

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 July 2002 (25.07.2002)

PCT

(10) International Publication Number
WO 02/057252 A2

(51) International Patent Classification⁷: C07D 333/34,
409/12, 307/64, A61K 31/381, 31/4535, 31/341, A61P
25/28

CASEBIER, David, Scott; 161 Ember Lane, Carlisle,
MA 01741 (US).

(21) International Application Number: PCT/US01/48375

(74) Agents: KODROFF, Cathy, A. et al.; Howson and How-
son, Spring House Corporate Center, P.O. Box 457, Spring
House, PA 19477 (US).

(22) International Filing Date:
11 December 2001 (11.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/255,105 13 December 2000 (13.12.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU,
ZA, ZM, ZW.

(71) Applicants: WYETH [US/US]; 5 Giralda Farms, Madi-
son, NJ 07940 (US). ARQULE, INC. [US/US]; 19 Presi-
dential Way, Woburn, MA 01801 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(72) Inventors: KREFT, Anthony, Frank; 43 Barley Court,
Langhorne, PA 19047 (US). COLE, Derek, Cecil; 4
Rheinlander Lane, New City, NY 10956 (US). WOLLER,
Kevin, Roger; 2 Amandrey Way, Ayer, MA 01741
(US). STOCK, Joseph, Raymond; 439 High Street,
Monroe, NY 10950 (US). DIAMANITIS, George; 106
Radtko Road, Randolph, NJ 07869 (US). KURBRAK,
Dennis, Michael; 3231 Gaul, Philadelphia, PA 19134
(US). KUTTERER, Kristina, Martha; 624 Cambell
Place, Westwood, NJ 07675 (US). MOORE, William,
Jay; 67 Ringold Street, Marlborough, MA 01752 (US).


Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/057252 A2

(54) Title: HETEROCYCLIC SULFONAMIDE INHIBITORS OF BETA AMYLOID PRODUCTION



(I)

(57) Abstract: Compounds of Formula (I), wherein R₁, R₂, R₃, R₄, R₅, R₆, T, W, X, Y and Z are as defined herein are provided, together with pharmaceutically acceptable salt, hy-
drates and/or prodrugs thereof. Methods of using these com-
pounds for inhibiting beta amyloid production and for treat-
ment of Alzheimer's Disease and Down's syndrome are de-
scribed.

HETEROCYCLIC SULFONAMIDE INHIBITORS OF BETA AMYLOID PRODUCTION

5

BACKGROUND OF THE INVENTION

This invention relates to inhibitors of beta amyloid production, which have utility in the treatment of Alzheimer's disease.

Alzheimer's Disease (AD) is the most common form of dementia (loss of
10 memory) in the elderly. The main pathological lesions of AD found in the brain consist of extracellular deposits of beta amyloid protein in the form of plaques and angiopathy and intracellular neurofibrillary tangles of aggregated hyperphosphorylated tau protein. Recent evidence has revealed that elevated beta amyloid levels in brain not only precede tau pathology but also correlate with
15 cognitive decline. Further suggesting a causative role for beta amyloid in AD, recent studies have shown that aggregated beta amyloid is toxic to neurons in cell culture.

Beta amyloid protein is composed mainly of 39-42 amino acid peptides and is produced from a larger precursor protein called amyloid precursor protein (APP) by the sequential action of the proteases beta and gamma secretase. Although rare, cases
20 of early onset AD have been attributed to genetic mutations in APP that lead to an overproduction of either total beta amyloid protein or its more aggregation-prone 42 amino acid isoform. Furthermore, people with Down's Syndrome possess an extra chromosome that contains the gene that encodes APP and thus have elevated beta amyloid levels and invariably develop AD later in life.

25 There continues to be an unmet need for compositions useful in inhibiting beta amyloid production and in the treatment of the effects of Alzheimer's Disease (AD).

SUMMARY OF THE INVENTION

The present invention provides heterocyclic sulfonamide derivatives of 2-
30 amino-1-alcohols and related homologs that have been found to specifically inhibit the production of beta amyloid protein from APP and to be capable of passing through the blood-brain barrier. These compounds are useful for the treatment of conditions

in which beta amyloid levels are elevated (e.g., AD, Down's Syndrome). Systemic administration of these compounds to subjects at risk of, or suffering from, these diseases lowers beta amyloid protein levels with subsequent reduction in the toxic beta amyloid aggregates in the brains of these patients.

5 In one aspect, the present invention provides a compound of Formula (I), as defined herein, pharmaceutically acceptable salts, hydrates, or prodrugs thereof. In one embodiment, the compounds of Formula (I) are thiophenesulfonamides. In another embodiment, the compounds of Formula (I) are furansulfonamides. Among the particularly desirable compounds are those having a halogen in the 5-position of
10 the heterocycle (e.g., 5-halo thiophenesulfonamides) and β -branches in the side chain of the primary alcohol.

In another aspect, the invention provides a pharmaceutical composition containing one or more compounds of Formula (I) and a physiologically compatible carrier.


15 In yet another aspect, the invention provides a method of inhibiting beta amyloid production in a subject by delivering a compound of Formula (I).

In still another aspect, the invention provides a method of treating Alzheimer's Disease (AD) in a subject by administering a compound of Formula (I) to the subject in an amount sufficient to alleviate the symptoms or progress of AD.

20 These and other aspects of the invention will be apparent to one of skill in the art upon reading of the following detailed description of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The invention consists of compounds of Formula (I), their
25 pharmaceutical formulations, and their use in modulating beta amyloid production in subjects at risk for, or suffering from, AD or other diseases resulting from elevated levels of beta amyloid protein in the brain. The compounds of Formula (I) include pharmaceutically acceptable salts and/or hydrates or prodrugs thereof, wherein:



(I)

R_1 and R_2 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, CF_3 , alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, phenyl, substituted phenyl, and

5 $(CH_2)_n(1,3)$ dioxane, where n is 2 to 5;

R_3 is selected from the group consisting of hydrogen, alkyl, and substituted alkyl;

R_4 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkylcycloalkyl, substituted alkylcycloalkyl, phenyl(substituted)alkyl, alkylOH, substituted alkylOH, alkylOBn, substituted alkylOBn, alkylpyridyl, substituted alkylpyridyl, alkylfuranyl, substituted alkylfuranyl, $CH(OH)$ phenyl, $CH(OH)$ substituted phenyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, N-substituted-piperidiny, piperidiny, substituted piperidiny, tetrahydrothiopyran, substituted tetrahydrothiopyran, 2-indane, substituted 2-indane, phenyl, substituted phenyl, alkylNHR₇, and substituted alkylNHR₇;

with the proviso that R_3 and R_4 are not both hydrogen;

R_7 is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, benzyl, substituted benzyl, alkylOH, substituted alkylOH, alkylSR₈, or substituted alkylSR₈;

R_8 is alkyl, substituted alkyl, benzyl, or substituted benzyl;

20 or R_3 and R_4 may be joined to form a ring;


R_5 is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CH_2 cycloalkyl, substituted CH_2 cycloalkyl, benzyl, substituted benzyl, and $CH_2CH_2QR_9$;

Q is O, NH or S;

25 R_9 is lower alkyl, substituted lower alkyl, phenyl, or substituted phenyl;

R_6 is selected from the group consisting of hydrogen, halogen and CF_3 ;

T is selected from the group consisting of



W, Y and Z are independently selected from the group consisting of C, CR₁₀ and N;

R₁₀ is selected from the group consisting of hydrogen and halogen, with the proviso that at least one of W, Y and Z must be C;

5 X is selected from the group consisting of O, S, SO₂, and NR₁₁;

R₁₁ is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, and substituted phenyl;

provided that when the compound contains one or more chiral centers, at least one of the chiral centers must be of S-stereochemistry.

10 The point of attachment of the W-X-Y-Z-C heterocyclic ring to the SO₂ group is not a limitation of the present invention. However, in one preferred embodiment, the ring is attached to the SO₂ group through a carbon-atom. However, the ring may be attached through O, S, or N heteroatoms.

The compounds of the invention may contain one or more asymmetric carbon
15 atoms and some of the compounds may contain one or more asymmetric (chiral) centers and may thus give rise to optical isomers and diastereomers. While shown without respect to stereochemistry in Formula (I), when the compounds of Formula (I) contain one or more chiral centers, at least one of the chiral centers is of S-stereochemistry. Most preferably, the carbon atom to which N, T, R₃ and R₄ are
20 attached is of S-stereochemistry. Thus, the invention includes such optical isomers and diastereomers; as well as the racemic and resolved, enantiomerically pure stereoisomers; as well as other mixtures of the R and S stereoisomers, and pharmaceutically acceptable salts, hydrates, and prodrugs thereof.

The term "alkyl" is used herein to refer to both straight- and branched-chain
25 saturated aliphatic hydrocarbon groups having one to ten carbon atoms, preferably one to eight carbon atoms and, most preferably, one to six carbon atoms; as used herein, the term "lower alkyl" refers to straight- and branched-chain saturated aliphatic hydrocarbon groups having one to six carbon atoms; "alkenyl" is intended to include both straight- and branched-chain alkyl group with at least one carbon-
30 carbon double bond and two to eight carbon atoms, preferably two to six carbon atoms; "alkynyl" group is intended to cover both straight- and branched-chain alkyl

groups with at least one carbon-carbon triple bond and two to eight carbon atoms, preferably two to six carbon atoms.

The terms "substituted alkyl", "substituted alkenyl", and "substituted alkynyl" refer to alkyl, alkenyl, and alkynyl as just described having from one to three
5 substituents selected from the group including halogen, CN, OH, NO₂, amino, aryl, heterocyclic, substituted aryl, substituted heterocyclic, alkoxy, substituted alkoxy, aryloxy, substituted alkyloxy, alkylcarbonyl, alkylcarboxy, alkylamino, arylthio. These substituents may be attached to any carbon of an alkyl, alkenyl, or alkynyl group provided that the attachment constitutes a stable chemical moiety.

10 The term "aryl" is used herein to refer to a carbocyclic aromatic system, which may be a single ring, or multiple aromatic rings fused or linked together as such that at least one part of the fused or linked rings forms the conjugated aromatic system. The aryl groups include, but are not limited to, phenyl, naphthyl, biphenyl, anthryl, tetrahydronaphthyl, phenanthryl, and indane.

15 The term "substituted aryl" refers to aryl as just defined having one to four substituents from the group including halogen, CN, OH, NO₂, amino, alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, aryloxy, substituted alkyloxy, alkylcarbonyl, alkylcarboxy, alkylamino, and arylthio.

The term "substituted benzyl" refers to a benzyl group, having substituted on
20 the benzene ring, one to five substituents from the group including halogen, CN, OH, NO₂, amino, alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, aryloxy, substituted alkyloxy, alkylcarbonyl, alkylcarboxy, alkylamino, and arylthio.

The term "heterocyclic" is used herein to describe a stable 4- to 7-membered monocyclic or a stable multicyclic heterocyclic ring which is saturated, partially
25 unsaturated, or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group including N, O, and S atoms. The N and S atoms may be oxidized. The heterocyclic ring also includes any multicyclic ring in which any of above defined heterocyclic rings is fused to an aryl ring. The heterocyclic ring may be attached at any heteroatom or carbon atom provided the
30 resultant structure is chemically stable. Such heterocyclic groups include, for example, tetrahydrofuran, piperidinyl, piperazinyl, 2-oxopiperidinyl, azepinyl, pyrrolidinyl, imidazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl,

isoxazolyl, morpholinyl, indolyl, quinolinyl, thienyl, furyl, benzofuranyl, benzothienyl, thiamorpholinyl, thiamorpholinyl sulfoxide, isoquinolinyl, and tetrahydrothiopyran.

The term "substituted heterocyclic" is used herein to describe the heterocyclic just defined having one to four substituents selected from the group which includes halogen, CN, OH, NO₂, amino, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, aryloxy, substituted aryloxy, alkyloxy, substituted alkyloxy, alkylcarbonyl, substituted alkylcarbonyl, alkylcarboxy, substituted alkylcarboxy, alkylamino, substituted alkylamino, arylthio, or substituted arylthio.

The term "substituted cycloalkyl" is used herein to describe a carbon-based ring having more than 3 carbon-atoms which forms a stable ring and having from one to five substituents selected from the group consisting of halogen, CN, OH, NO₂, amino, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, alkoxy, aryloxy, substituted alkyloxy, alkylcarbonyl, alkylcarboxy, alkylamino, substituted alkylamino, arylthio, heterocyclic, substituted heterocyclic, aminoalkyl, and substituted aminoalkyl.

Where the terms "substituted alkylcycloalkyl", "substituted alkylOBn", "substituted alkylpyridyl", "substituted alkylfuryl", "substituted alkyl NHR₇", "substituted alkylOH", and "substituted alkylSR₈" are recited, the substitution may occur at the alkyl group or on the corresponding base compound.

As used in the definition of the R₄ group, an N-substituted piperidinyl group may be defined as are the substituted heterocyclic groups. Among particularly desirable substituents are N-alkyl-, N-aryl-, N-acyl-, and N-sulfonyl piperidinyl groups. One particularly suitable N-acyl-piperidinyl group is N-t-butyloxycarbonyl (BOC)-piperidine. However, other suitable substituents can be readily identified by one of skill in the art.

The term "alkoxy" is used herein to refer to the OR group, where R is alkyl or substituted alkyl. The term "aryloxy" is used herein to refer to the OR group, where R is aryl or substituted aryl. The term "alkylcarbonyl" is used herein to refer to the RCO group, where R is alkyl or substituted alkyl. The term "alkylcarboxy" is used herein to refer to the COOR group, where R is alkyl or substituted alkyl. The term

"aminoalkyl" refers to both secondary and tertiary amines wherein the alkyl or substituted alkyl groups, containing one to eight carbon atoms, which may be either same or different and the point of attachment is on the nitrogen atom.

The term "halogen" refers to Cl, Br, F, or I.

- 5 The term "ring" structure, e.g., when R₃ and R₄ may form a ring structure, includes a monocyclic structure, a bridged cyclo structure, and fused cyclo structures, unless the type of ring structure is otherwise specified.

- The compounds of the present invention can be used in the form of salts derived from pharmaceutically or physiologically acceptable acids or bases. These salts include, but are not limited to, the following salts with organic and inorganic acids such as acetic, lactic, citric, tartaric, succinic, fumaric, maleic, malonic, mandelic, mallic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, toluenesulfonic and similarly known acceptable acids, and mixtures thereof. Other salts include salts with alkali metals or alkaline earth metals, such as sodium (e.g., sodium hydroxide), potassium (e.g., potassium hydroxide), calcium or magnesium.
- 10
- 15

- These salts, as well as other compounds of the invention may be in the form of esters, carbamates and other conventional "pro-drug" forms, which, when administered in such form, convert to the active moiety *in vivo*. In a currently preferred embodiment, the prodrugs are esters. See, e.g., B. Testa and J. Caldwell, "Prodrugs Revisited: The "Ad Hoc" Approach as a Complement to Ligand Design", Medicinal Research Reviews, 16(3):233-241, ed., John Wiley & Sons (1996).
- 20

- In one particularly desirable embodiment, the compounds of Formula (I) are thiophenesulfonamides, and more desirably, 5-halo thiophenesulfonamides, and most desirably, 5-halo thiophene sulfonamides with β -branches in the side chain of a primary alcohol. Thus, with respect to Formula (I), the compound of the invention desirably has a structure in which X is S, W is C (or CR₁₀), Y is C (or CR₁₀) and Z is C (or CR₁₀), and the sulfonamide is attached to C2 of the thiophene ring. More desirably, X is S, W is C (or CR₁₀), Y is C (or CR₁₀), Z is C (or CR₁₀) and R₆ is a halogen. Most desirably, X is S, X is C, W is C, Y is C, Z is C, R₆ is a halogen, and T is C(OH)R₁R₂, where R₁ and R₂ are hydrogen, R₃ is H, R₄ is a lower alkyl of S-stereochemistry, and R₅ is H. In preliminary screening assays *in vitro* and *in vivo*,
- 25
- 30

compounds of these structures have been found to have unexpectedly good beta-amyloid inhibitory activity, and in many cases, better activity than compounds of Formula (I) having other heterocycles (e.g., furans, where X is O). However, other such compounds of Formula (I) are also useful for the purposes described herein.

5 For example, in another embodiment, the compounds of Formula (I) are furansulfonamides, in which X is O, W is C, Y is C, and Z is C. In one particularly desirable embodiment, the furansulfonamides of Formula (I) are further characterized by β -branches in the side chain of a primary alcohol. Thus, with respect to Formula (I), in these compounds T is C(OH)R₁R₂, in which R₁ and R₂ are hydrogen, R₃ is H,
10 R₄ is a lower alkyl of S-stereochemistry, R₅ is H and R₆ is halogen.

 In still another embodiment, the compounds of Formula (I) are characterized by being sulfonamides of Formula (I), which have β -branches in the side chain of the primary alcohol group. Thus, with respect to Formula (I), in these compounds T is C(OH)R₁R₂, R₁ and R₂ are hydrogen, R₃ is H, R₄ is a lower alkyl of S-
15 stereochemistry, and R₅ is H.

 These and the other compounds of the invention can be prepared following the Schemes illustrated below.


Synthesis

20 The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention can be prepared using the methods described below, together with synthetic methods known in the synthetic organic arts or variations of these methods by one skilled in the art. (See, generally, *Comprehensive Organic Synthesis*,
25 "Selectivity, Strategy & Efficiency in Modern Organic Chemistry", ed., I. Fleming, Pergamon Press, New York (1991); *Comprehensive Organic Chemistry*, "The Synthesis and Reactions of Organic Compounds", ed. J.F. Stoddard, Pergamon Press, New York (1979)). Preferred methods include, but are not limited to, those outlined below.

30 A first method of preparation consists of reaction of a 2-aminoalcohol II with the appropriate sulfonyl halide in the presence of a base such as triethylamine (TEA) and in a suitable solvent to afford compounds of Formula III. For compounds where

- R_2 and R_1 are hydrogen, oxidation of the N-sulfonyl primary alcohol with pyridinium chlorochromate (PCC) or under Swern conditions then affords the corresponding aldehyde IV which can be reacted with Grignard reagents ($RMgX$, where R is an organic radical and X is a halogen) to afford the secondary alcohols V as a mixture of diastereomers which can be separated by high performance liquid chromatography (HPLC) (Scheme 1).

Scheme 1




- A second method of preparation involves reaction of an α -amino acid or ester IX with the appropriate sulfonyl halide in the presence of a base such as triethylamine and in a suitable solvent to afford compounds of Formula X (Scheme 2). The intermediate N-sulfonyl acid X ($R_x=\text{H}$) can be converted to the corresponding primary alcohol VIII ($R_1=R_2=\text{H}$) utilizing standard methodology such as LiAlH_4 , B_2H_6 or cyanuric chloride/ NaBH_4 . The intermediate N-sulfonyl ester X ($R_x=\text{alkyl}$, Bn) can also be reduced to the corresponding primary alcohol VIII ($R_1=R_2=\text{H}$) utilizing standard methodology such as LiAlH_4 . Alternatively, the intermediate N-sulfonyl ester X ($R_x=\text{alkyl}$, Bn) can be converted to the aldehyde IV with DiBAL. Finally, the intermediate N-sulfonyl ester X ($R_x=\text{alkyl}$, Bn) can be reacted with 2 equivalents of Grignard reagent to afford the tertiary alcohols III with $R_1=R_2$. Alternatively, for tertiary alcohols III with R_1 not equal to R_2 , the corresponding

Weinreb amide (see Scheme 10) of the N-sulfonyl acid can be prepared and subsequently reacted with R_1MgX and R_2MgX . For compounds of formula X ($R_x=H$) that have an asymmetric center at the α -amino acid carbon, the pure enantiomers can be obtained by standard resolution procedures employing

5 recrystallization of salts formed with various chiral bases.

Scheme 2




In a variation of the second method to prepare the primary alcohols, an α -amino acid or ester (or N-protected derivative thereof) VI is first converted to the corresponding primary 2-aminoalcohol VII (using the methodology outlined in the previous

10 paragraph), which is subsequently, after deprotection (if necessary), reacted with the appropriate sulfonyl halide (Scheme 3) to afford compounds of Formula VIII. For preparation of compounds derived from unnatural α -amino acids containing beta branching in the amino acid side chain, a method of preparation based on the work of Hruby (*Tet. Lett.* 38: 5135-5138 (1997)) is outlined in Scheme 4. This route entails

15 formation of the α,β -unsaturated amide XII of the Evans chiral auxiliary from an α,β -unsaturated acid XI, followed by conjugate addition of an organocuprate,


trapping of the resulting enolate anion XIII with NBS, displacement of the bromide XIV with azide anion (provided by tetramethylguanidinium azide (TMGA)) to afford XV, followed by reduction to the 2-amino alcohol and subsequent sulfonylation to afford the target compound XVI. In Schemes 1 through 4, R_3 is H.

Scheme 3




5

Scheme 4



For the preparation of N-alkylated sulfonamides VIII (R_5 =alkyl etc.), the sulfonamide ester XVII can be N-alkylated by either treatment with a suitable base such as potassium carbonate followed by the alkylating agent R_5X or by employing Mitsunobu conditions ($R_5OH/DEAD$, TPP). $LiBH_4$ reduction of the N-alkylated sulfonamide ester affords the N-alkylated sulfonamide in the primary alcohol series VIII (Scheme 5). These primary alcohols VIII can be converted to the secondary alcohols V or aldehyde IV series by chemistry that has been outlined above. Alternatively, the N-alkylated sulfonamide esters, or their corresponding Weinreb amides, can be treated with Grignard reagents to afford the N-alkylated tertiary alcohols III.

Scheme 5



When the heterocycle attached to the sulfonamide in the above alcohols is thiophene, the corresponding sulfone derivative XIX may be obtained by oxidation of the thiophene compound XVIII with MCPBA (Scheme 6).


Scheme 6



An alternate preparation of sulfonamides derived from unnatural 2-aminoalcohols utilizes the Bucherer modification of the Strecker α -amino acid synthesis (Scheme 7). In this route, an aldehyde XX is reacted with cyanide anion


and ammonium carbonate to afford the hydantoin XXI, which is hydrolyzed to the α -amino acid XXII. This compound is then reduced to XXIII and sulfonylated to afford the desired compounds of Formula XXIV.

Scheme 7



- 5 For sulfonamides derived from 2-aminoalcohols containing an N or O heteroatom in the side chain, a route has been devised starting from D-serine (Scheme 8). In this route, D-serine XXV is first sulfonylated to XXVI and


Scheme 8



subsequently converted to the ketone XXVII, which is reductively aminated to the target compounds of Formula XXVIII.


For sulfonamides derived from 2-aminoalcohols in the secondary alcohol series with $R_1=H$ and $R_2=CF_3$ (compound XXIX), a method of preparation has been devised that is outlined in Scheme 9 starting from the aldehyde IV (prepared as in Scheme 1).

Scheme 9



As has been mentioned in the section concerning Scheme 1, the preparation of sulfonamides derived from 2-aminoalcohols in the secondary alcohol series V results in the formation of a diastereomeric mixture. An alternate method of preparation of these compounds that results in the production of a pure diastereomer is outlined in Scheme 10 for compounds derived from L-isoleucine. This method, which utilizes chemistry previously employed by Roux (*Tetrahedron* 50: 5345-5360 (1994)), consists of addition of Grignard reagents to the Weinreb amide XXX (derived from the requisite α -amino acid) followed by stereospecific reduction of the ketone XXXI to afford a single diastereomeric N-protected 2-amino alcohol XXXII. Deprotection of this compound followed by reaction with sulfonyl chlorides affords the pure diastereomeric sulfonamide secondary alcohols of Formula XXXIII.

Scheme 10



When the heterocycle attached to the sulfonamide in the above alcohols is thiophene, the corresponding 5-iodo and 5-fluoro-thiophene derivatives may be obtained by conversion of the 5-bromo-thiophene derivative XXXIV (obtained as in


5 Scheme 1) to a 5-trialkyltin-thiophene intermediate XXXV which can be converted to either the 5-iodo-thiophene (XXXVII) by treatment with sodium iodide and chloramine T or the 5-fluoro-thiophene analog (XXXVI) by treatment with SELECTFLUOR™ (Aldrich Chemical Co) (Scheme 11).

Sulfonamides derived from cyclohexylglycinol substituted by alkoxy and


10 amino groups at the 4 position of the cyclohexane ring can be prepared according to the methods described herein (Scheme 12). This route entails initial hydrogenation of 4-L-hydroxyphenylglycine XXXVIII, followed by sulfonylation, reduction of the carboxylic acid with diborane and formation of the N,O-acetonide XXXIX. The 4-hydroxy acetonide XXXIX is then O-alkylated using sodium hydride and an

15 alkylating agent such as an alkyl or benzyl bromide. This is followed by removal of the protecting group by treatment with aqueous acid to afford the 4-ether derivatives of Formula XXXX. Alternatively, the 4-hydroxy acetonide XXXIX can be oxidized to the 4-ketone which can be reductively aminated and deprotected to afford the corresponding 4-amino analogs of Formula XXXXI.

Scheme 11




Scheme 12



Another method of preparing chirally pure N-sulfonyl 2-amino alcohols derived from α -amino acids is outlined in Scheme 13. This method involves construction of an Evans oxazolidone chiral auxiliary XXXXIII from XXXXII, which is then converted to the corresponding enolate and electrophilically aminated with


5 trisyl azide to afford the key intermediate XXXXIV (*J. Am. Chem. Soc.* 109: 6881-6883 (1987)). The azide intermediate XXXXIV is then hydrolyzed to the α -azido acid XXXXV and reduced to the chirally pure α -amino acid XXXXVI which can be converted to the corresponding N-sulfonyl 2-amino alcohols by methods previously described above (e.g. Scheme 2).

10




Finally, chirally pure α -amino acids XXXXVI, one of the possible synthetic precursors of chiral N-sulfonyl 2-amino alcohols as mentioned above, can also be prepared utilizing an asymmetric variant of the Strecker α -amino acid synthesis as

15 outlined in Scheme 14 (*J. Org. Chem.* 54:1055-1062 (1989)).



Oximes XXXXXIV can be derived from the corresponding aldehydes IV by standard methodology as depicted in Scheme 15.



10

Methods of Use

Compounds of Formula (I) are inhibitors of beta amyloid production. In preliminary studies using protease specific assays, exemplary compounds of Formula (I) have been shown to exhibit specific inhibition with respect to protease activity. Thus, the compounds of the present invention are useful for treatment and prevention of a variety of conditions in which modulation of beta amyloid levels provides a therapeutic benefit. Such conditions include, e.g., amyloid angiopathy, cerebral amyloid angiopathy, systemic amyloidosis, Alzheimer's Disease (AD), hereditary cerebral hemorrhage with amyloidosis of the Dutch type, inclusion body myositis, Down's syndrome, among others.

20

In addition, the compounds of Formula (I) may be utilized in generating reagents useful in diagnosis of conditions associated with abnormal levels of beta

amyloid. For example, the compounds of Formula (I) may be used to generate antibodies, which would be useful in a variety of diagnostic assays. Methods for generating monoclonal, polyclonal, recombinant, and synthetic antibodies or fragments thereof, are well known to those of skill in the art. (See, e.g., E. Mark and
5 Padlin, "Humanization of Monoclonal Antibodies", Chapter 4, *The Handbook of Experimental Pharmacology*, Vol. 113, The Pharmacology of Monoclonal Antibodies, Springer-Verlag (June, 1994); Kohler and Milstein and the many known modifications thereof; PCT Patent Application No. PCT/GB85/00392; British Patent Application Publication No. GB2188638A; Amit et al., *Science*, **233**:747-753 (1986);
10 Queen et al., *Proc. Nat'l. Acad. Sci. USA*, **86**:10029-10033 (1989); International Patent Publication No. WO90/07861; and Riechmann et al., *Nature*, **332**:323-327 (1988); Huse et al., *Science*, **246**:1275-1281 (1988)). Alternatively, the compounds of Formula (I) may themselves be used in such diagnostic assays. Regardless of the reagent selected (e.g., antibody or compound of Formula (I)), suitable diagnostic
15 formats including, e.g., radioimmunoassays and enzyme-linked immunosorbent assays (ELISAs), are well known to those of skill in the art and are not a limitation on this embodiment of the invention.

The beta amyloid inhibitory activity of many of the compounds of the present invention has been determined using the Repressor Release Assay (RRA). See, Table
20 23 below. A compound is considered active in RRA if it leads to at least a 1.5 fold increase in luciferase activity at 20 μ M and is non-toxic.

Additionally, cellular, cell-free and in vivo screening methods to detect inhibitors of beta amyloid production are known in the art. Such assays may include radioimmunoassays and enzyme-linked immunosorbent assay (ELISA), among
25 others. See, e.g., P.D. Mehta, et al., *Techniques in Diagnostic Pathology*, vol. 2, eds., Bullock et al, Academic Press, Boston, pages 99-112 (1991), International Patent Publication No. WO 98/22493, European Patent No. 0652009, US Patent No. 5,703,129 and US Patent No. 5,593,846. Selection of an appropriate in vitro or in vivo screening assay is not a limitation of the present invention.

30

Pharmaceutical Formulation

The compounds of this invention may be administered to a subject by any desirable route, taking into consideration the specific condition for which it has been selected. By subject is meant any suitable mammal, including humans, domestic
5 animals (e.g., canines and felines), and livestock, which have been recognized as having or at risk of having one or more of the conditions for which modulation of beta amyloid levels is desirable. Thus, the compounds of the invention are useful for treatment and/or prevention of a number of human and veterinary conditions. As used
10 herein, "prevention" encompasses prevention of symptoms in a subject who has been identified as at risk for the condition, but has not yet been diagnosed with the same and/or who has not yet presented any symptoms thereof.

These compounds may be delivered or administered by any suitable route of delivery, e.g., oral, intravenous, subcutaneous, intramuscular, sublingual, intracranial, epidural, intratracheal, rectal, vaginal, among others. Most desirably, the compounds
15 are delivered orally or by a suitable parenteral route. The compounds may be formulated in combination with conventional pharmaceutical carriers that are physiologically compatible. Optionally, one or more of the compounds of the invention may be mixed with other active agents.

Suitable physiologically compatible carriers may be readily selected by one of
20 skill in the art. For example, suitable solid carriers include, among others, one or more substances which may also act as lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents or an encapsulating material. In powders, the carrier is a finely divided solid, which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is
25 mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, starch, sugars (including, e.g., lactose and sucrose), dicalcium phosphate, cellulose (including, e.g., microcrystalline cellulose, methyl cellulose, sodium
30 carboxymethyl cellulose), and kaolin.

Liquid carriers may be used in preparing solutions, suspensions, emulsions, syrups and elixirs. The active ingredient of this invention can be dissolved or

suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, suspending agents, thickening agents, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives, and oils (e.g., fractionated coconut oil, arachis oil, corn oil, peanut oil, and sesame oil). For parenteral administration the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration.

Optionally, additives customarily employed in the preparation of pharmaceutical compositions may be included in the compositions of the invention. Such components include, e.g., sweeteners or other flavoring agents, coloring agents, preservatives, and antioxidants, e.g., vitamin E, ascorbic acid, BHT and BHA.

Liquid pharmaceutical compositions that are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Oral administration may be either liquid or solid composition form.

Preferably the pharmaceutical composition is in unit dosage form, e.g. as tablets or capsules. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage forms can be packaged compositions, for example packeted powders, vials, ampoules, prefilled syringes or sachets containing liquids. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

As described herein, a therapeutically or prophylactically useful amount of a compound of the invention is that amount of a compound which alleviates the symptoms of the disease, e.g., AD, or which prevents the onset of symptoms, or the onset of more severe symptoms. Generally, an individual dose (i.e., per unit, e.g., tablet) of a compound of the invention may be in the range from about 1 $\mu\text{g/kg}$ to

about 10 g/kg, more preferably 10 mg/kg to about 5 g/kg, and most preferably about 1 mg/kg to about 200 mg/kg. Desirably, these amounts are provided on a daily basis.


However, the dosage to be used in the treatment or prevention of a specific cognitive deficit or other condition may be subjectively determined by the attending physician.

- 5 The variables involved include the specific cognitive deficit and the size, age and response pattern of the patient. For example, based upon the activity profile and potency of the compounds of this invention, a starting dose of about 10 mg per day with gradual increase in the daily dose to about 200 mg per day may provide the desired dosage level in the human.
- 10 Alternatively, the use of sustained delivery devices may be desirable, in order to avoid the necessity for the patient to take medications on a daily basis. "Sustained delivery" is defined as delaying the release of an active agent, i.e., a compound of the invention, until after placement in a delivery environment, followed by a sustained release of the agent at a later time. Those of skill in the art know suitable sustained
- 15 delivery devices. Examples of suitable sustained delivery devices include, e.g., hydrogels (see, e.g., US Patent Nos. 5,266,325; 4,959,217; and 5,292,515), an osmotic pump, such as described by Alza (US Patent No. 4,295,987 and US Patent No. 5,273,752) or Merck (European Patent No. 314,206), among others; hydrophobic membrane materials, such as ethylenemethacrylate (EMA) and ethylenevinylacetate
- 20 (EVA); bioresorbable polymer systems (see, e.g., International Patent Publication No. WO 98/44964, Bioxid and Cellomeda; US Patent No. 5,756,127 and US Patent No. 5,854,388); other bioresorbable implant devices have been described as being composed of, for example, polyesters, polyanhydrides, or lactic acid/glycolic acid copolymers (see, e.g., US Patent No. 5,817,343 (Alkermes Inc.)). For use in such
- 25 sustained delivery devices, the compounds of the invention may be formulated as described herein.

EXAMPLES


The following examples are provided to illustrate the production and activity of representative compounds of the invention and to illustrate their performance in a screening assay. One skilled in the art will appreciate that although specific reagents and conditions are outlined in the following examples, these reagents and conditions are not a limitation on the present invention.

Example 1

3-Bromo-5-chloro-N-[(1S, 2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide


To a solution of (S)-+- isoleucinol (23 mg, 0.2 mmol) in THF (3 mL) was added triethylamine (46 μ L, 0.24 mmol) and 3-bromo-5-chlorothiophene-2-sulfonyl chloride (59.2 mg, 0.2 mmol). The solution was stirred for 8 – 16 h, then concentrated. The residue was dissolved in MeOH (1.5 mL) and purified by semi-preparative RP-HPLC¹ to give Example 1 (20.3 mg).

The following compounds (Examples 1-7, Table 1) were prepared using 3-bromo-5-chlorothiophene-2-sulfonyl chloride, 5-bromothiophene-2-sulfonyl chloride, 3-bromo-2-chlorothiophene-5-sulfonyl chloride, 5-chlorothiophene-2-sulfonyl chloride, 2,5-dichlorothiophene-3-sulfonyl chloride, 2,3-dichlorothiophene-5-sulfonyl chloride, and 2-thiophenesulfonyl chloride and following the procedure outlined in Example 1.

**Table 1 (LCMS² Data: Molecular ion and retention time)**

RSO ₂ Cl	(S)-(+)-isoleucinol
3-bromo-5-chlorothiophene-2-sulfonyl chloride	Example 1 (377 M+H); 3.25 min
5-bromothiophene-2-sulfonyl chloride	Example 2 (344 M+H); 3.01 min
3-bromo-2-chlorothiophene-5-sulfonyl chloride	Example 3 (378 M+H); 3.35 min
5-chlorothiophene-2-sulfonyl chloride	Example 4 (298 M+H); 2.97 min
2,5-dichlorothiophene-3-sulfonyl chloride	Example 5 (332 M+H); 3.18 min
2,3-dichlorothiophene-5-sulfonyl chloride	Example 6 (332 M+H); 3.33 min
2-thiophenesulfonyl chloride	Example 7 (264 M+H); 2.35 min

5


Example 8**5-Chloro-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide**

10

To a solution of L-valinol (25.8 mg, 0.25 mmol) in THF (3 mL) was added triethylamine (58 μ L, 0.3 mmol) and 5-chlorothiophene-2-sulfonyl chloride (54 mg, 0.25 mmol). The solution was stirred for 8 to 16 h, then concentrated. The residue

was dissolved in MeOH (1.5 mL) and purified by semi-preparative RP-HPLC¹ to give **Example 8** (19.5 mg).

The following compounds (**Examples 8-10**, **Table 2**) were prepared using 5-thiophene-2-sulfonyl chloride and 5-bromothiophenesulfonyl chloride with L-valinol and D-valinol and following the procedure outlined in **Example 8**.




10

Table 2 (LCMS² Data: Molecular ion and retention time)

	RSO ₂ Cl	
Amine	5-chlorothiophene-2-sulfonyl chloride	5-bromothiophene-2-sulfonyl chloride
L-valinol	Example 8 (284 M+H); 2.70 min	Example 9 (330 M+H); 2.75 min
D-valinol		Example 10 (330 M+H); 2.75 min

Example 11


4,5-Dibromo-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide



15

To a solution of (S)-(+)-2-amino-3-methyl-1-butanol (20.6 mg, 0.2 mmol) in THF (3 mL) was added triethylamine (46 μ L, 0.24 mmol) and 4,5-dibromothiophene-2-sulfonyl chloride (68 mg, 0.2 mmol). The solution was stirred for 8 to 16 h, the

solvent was removed and residue purified by RP-HPLC¹ to give **Example 11** (49.6 mg).




5

Table 3 (LCMS² Data: Molecular ion and retention time)

	NH₂CH(R₁)CH₂OH
RSO₂Cl	(S)-(+)-2-amino-3-methyl-1-butanol
4,5-dibromothiophene-2-sulfonyl chloride	Example 11 (408 M+H); 3.22 min

Example 12

10 **5-Chloro-N-[(1S)-1-cyclohexyl-2-hydroxyethyl]thiophene-2-sulfonamide**



A. Part 1

To a solution of L-cyclohexyl-glycine (48.5 mg, 0.25 mmol) in THF
 15 (2 mL) was added lithium aluminum hydride (1 M solution in THF) (0.8 mL, 0.8 mmol) and the solution heated at 60°C for 4 h. The solution was stirred at 25° C for 8 to 16 hours. The reaction was quenched by addition of water (45 µL), 15% aqueous sodium hydroxide (45 µL) and water (105 µL) with vigorous stirring between each addition. The mixture was then filtered and concentrated.

20

B. Part 2

To a solution of the residue from Part 1 in THF (3 mL) was added triethylamine (69 μ L, 0.50 mmol) and 5-chlorothiophene-2-sulfonyl chloride (54.3 mg, 0.25 mmol). The solution was stirred for 8 to 16 h, the solvent was removed and residue purified by RP-HPLC¹ to give **Example 12** (25.9 mg).

The following compounds (**Examples 12 - 17, Table 4**) were prepared using 5-chlorothiophene-2-sulfonyl chloride, and 5-bromothiophene-2-sulfonyl chloride with L-cyclohexylglycine, β -methyl-DL-phenylalanine, and L-allo-isoleucine and following the procedure outlined in Example 12.





Table 4 (LCMS² Data: Molecular ion and retention time)

Amino acid	RSO ₂ Cl	
	5-chlorothiophene-2-sulfonyl chloride	5-bromothiophene-2-sulfonyl chloride
L-cyclohexylglycine	Example 12 (324 M+H); 3.07 min	Example 13 (370 M+H); 3.10 min
beta-methyl-DL-phenylalanine	Example 14 (346 M+H); 3.05 min	Example 15 (392 M+H); 3.08 min
L-allo-isoleucine	Example 16 (298 M+H); 2.78 min	Example 17 (344 M+H); 2.82 min

Example 18**5-Bromo-N-[(1S, 2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
1,1-dioxide**

5

A. Part 1.


To a solution of (S)-+isoleucinol (58.6 mg, 0.5 mmol) in DCM (5 mL) was added triethylamine (210 μ L, 1.5 mmol) and 5-bromothiophene-2-sulfonyl chloride (130.8 mg, 0.5 mmol). The solution was stirred for 8 to 16 h, then concentrated.

10

B. Part 2.

The residue from Part 2 (0.5 mmol) was dissolved in dichloromethane (3 mL) and *meta*-chloroperbenzoic acid (2.5 mmol) was added. The solution was stirred for 8 to 16 h, the solvent was removed and residue purified by RP-HPLC to give **Example 18** (4.3 mg). LCMS² Data: Molecular ion and retention time, 375.9 M+H; 3.37 min.

15

Example 19**5-Chloro-N-[1-(hydroxymethyl)-2,3-dimethylpentyl]thiophene-2-sulfonamide****A. Part 1**

5 To a solution of sodium cyanide (735.15 mg, 15 mmol) and ammonium carbonate (1.92 g, 20 mmol) in EtOH/H₂O (1:1, 35 mL) was added 2,3 dimethylpentanal (570.95 mg, 5 mmol). The solution was heated at 50°C for 20 h, then concentrated.

B. Part 2

10 The residue from Part 1 (5 mmol) was dissolved in 35 mL of a 3N sodium hydroxide solution and heated at 95° C for 22 h. Stirring was continued for an additional 8 to 16 h, then the solvent was removed.

C. Part 3

15 To the residue from Part 2 (2.5 mmol) in THF (10 mL) was added lithium aluminum hydride (1 M solution in THF) (5 mL, 5 mmol) and the solution heated at 60°C for 4 h. The solution was stirred at 25° C for 8 to 16 h. The reaction was quenched by addition of water (285 µL), 15% aqueous sodium hydroxide (285 µL), and water (665 µL) with vigorous stirring between each addition. The mixture was then filtered and concentrated.

D. Part 4

20 To the residue from Part 3 (0.5 mmol) in THF (5 mL) was added triethylamine (83.7 µL, 0.6 mmol) and 5-chlorothiophene-2-sulfonyl chloride (108.54 mg, 0.5 mmol). The solution was stirred for 8 to 16 h, the solvent was removed and residue purified by RP-HPLC¹ to give **Example 19** (46.1 mg).

The following compounds (Examples 19 - 24, Table 5) were prepared using 2,3 dimethylpentanal, 2-methylvaleraldehyde, 2-ethylhexanal, 2,4,6-trimethyl-3-cyclohexene-1-carboxaldehyde, 1,2,3,6-tetrahydro-benzaldehyde, cyclopentylmethanal, and following the procedure outlined in Example 19.

5





Table 5 (LCMS² Data: Molecular ion and retention time)

	R''SO ₂ Cl
Aldehyde	5-chlorothiophene-2-sulfonyl chloride
2,3 dimethylpentanal	Example 19 (326 M+H); 3.47 min
2-methylvaleraldehyde	Example 20 (312 M+H); 3.25 min
2-ethylhexanal	Example 21 (340 M+H); 3.74 min
2,4,6-trimethyl-3-cyclohexene-1-carboxaldehyde	Example 22 (364 M+H); 3.76 min
1,2,3,6-tetrahydrobenzaldehyde	Example 23 (322 M+H); 3.11 min
Cyclopentylmethanal	Example 24 (310 M+H); 3.07 min

10

Example 25

5-Bromo-N-[(1S)-1-(hydroxymethyl)-1,2-dimethylpropyl]thiophene-2-sulfonamide



5

A. Part 1

To a solution of (S)- α -methyl valine (131 mg, 1 mmol) in THF (5 mL) was added lithium aluminum hydride (1 M solution in THF) (2 mL, 2 mmol) and the solution was heated at 60°C for 4 h. The solution was stirred at 25°C for 8 to 16 h.

10 The reaction was quenched by addition of water (114 μ L), 15% aqueous sodium hydroxide (114 μ L), and water (266 μ L) with vigorous stirring between each addition. The mixture was then filtered and concentrated.

B. Part 2

To the residue from Part 1 (0.5 mmol) in THF (2 mL) was added triethylamine (83.7 μ L, 0.6 mmol) and 5-bromothiophene-2-sulfonyl chloride (130.8 mg, 0.5 mmol). The solution was stirred for 8 to 16 h, the solvent was removed and residue purified by RP-HPLC¹ to give Example 25 (50.8 mg).

The following compounds (Examples 25-26, Table 6) were prepared using 5-bromothiophene-2-sulfonyl chloride and 5-chlorothiophene-2-sulfonyl chloride and following the procedure outlined in Example 25.

20





Table 6 (LCMS² Data: Molecular ion and retention time)

RSO ₂ Cl	S- α -methyl-valine
5-bromothiophene-2-sulfonyl chloride	Example 25 (344 M+H); 2.97 min
5-chlorothiophene-2-sulfonyl chloride	Example 26 (298 M+H); 2.92 min

Example 27

5 **5-Chloro-N-[(1S,2R)-1-(hydroxymethyl)-2,4-dimethylpentyl]thiophene-2-sulfonamide**

**A. Part 1**

A solution of 4-methyl-2-pentenoic acid (7.6 mL, 40 mmol) in THF (100 mL) was cooled to -78°C. Triethylamine (5.85 mL, 42 mmol) and
 10 trimethylacetyl chloride (pivaloyl chloride) (5.17 mL, 42 mmol) were added via syringe in that order. The dry ice bath was replaced with an ice bath and the reaction stirred at 0°C for 1 h, then the reaction was recooled to -78°C.

In a separate flask (R)-(+)-4-benzyl-2-oxazolidinone (7.0 g, 40 mmol) was dissolved in THF (100 mL) and cooled to -78°C, then n-butyl lithium (1.6 M, 25
 15 mL) was added via syringe. The mixture was stirred for 20 min then the above reaction mixture added by removing the septum and pouring quickly from one flask to the other (Note: attempts to transfer reaction mixture via cannula failed due to the suspended trimethylammonium chloride in the mixture).

The resulting mixture was stirred at -78°C for 30 min then allowed to
 20 warm to 25° C for 1 to 2 h before quenching with saturated aqueous NH₄Cl solution

(100 mL). Volatiles were removed on the rotary evaporator and the aqueous slurry was diluted with water (200 mL) and extracted with ethyl acetate (2 x 200 mL). The combined organic phase was dried over anhydrous MgSO_4 , filtered and concentrated. The product may crystallize out of solution and be of high purity. If purification is required the crude product may be purified by flash chromatography using 20 to 30% ethyl acetate in hexane.

B. Part 2

To a copper (I) bromide/dimethyl sulfide complex (246 mg, 1.2 mmol) in THF/DMS (2:1, 15 mL), cooled to -40°C , was added methyl magnesium bromide (2.4 mL, 1 M solution in THF, 2.4 mmol). The solution was allowed to stir for 10 min while warming to -15°C . The mixture was recooled to -40°C and the product from Part 1 (245 mg, 1 mmol) in THF (6 mL) was added. The solution was stirred at 25°C for 8 to 16 h. The solution was recooled to -78°C and N-bromosuccinimide (356 mg, 2 mmol) in THF (2 mL) was added. The solution was allowed to warm to 0°C and shaken at 0°C for 3 h. The reaction was quenched with a 1:1 solution of saturated ammonium carbonate and 0.5 N potassium bisulfate (5 mL). The organic phase was decanted off and concentrated.

C. Part 3


To the product from Part 3 dissolved in acetonitrile (5 mL) was added tetramethylguanidine azide (0.6 mL, 4 mmol). The solution was stirred for 72 to 120 h. The solution was concentrated to dryness, redissolved in CH_2Cl_2 and 1 N HCl (2 mL) was added. The layers were separated and the organic layer was filtered through a pad of silica gel washed with CH_2Cl_2 (5 mL) and concentrated.

D. Part 4

To the product from Part 3 (131 mg, 1 mmol) in THF (5 mL) at 0°C was added lithium aluminum hydride (1 M solution in THF) (2 mL, 2 mmol) and the solution stirred at 25°C for 4 h. The reaction was quenched by addition of water (114 μL), 15% aqueous sodium hydroxide (114 μL), and water (266 μL) with vigorous stirring between each addition. The mixture was then filtered and concentrated.

E. Part 5

To the residue from Part 4 (0.5 mmol) in THF (2 mL) was added triethylamine (83.7 μ L, 0.6 mmol) and 5-chlorothiophene-2-sulfonyl chloride (108 mg, 0.5 mmol). The solution was stirred for 8 to 16 h, the solvent was removed and residue purified as described for Example 1 to give 50.8 mg.



R= Me, Et, n-Pr, i-Pr, hexyl, phenyl
biphenyl, 3-pyridyl, 2-furyl

R'MgX= Me, Et, i-Bu, hexyl, phenyl, 4-MeOPh

The following compounds (Examples 27-55, Table 7) were prepared using 5-chlorothiophene-2-sulfonyl chloride with crotonic acid, 2-pentenoic acid, 2-hexenoic acid, 2-octenoic acid, cinnamic acid, furylacrylic acid, 4-methyl-2-pentenoic acid, and 4-phenylcinnamic acid and methyl, ethyl, isobutyl, 4-methoxyphenyl, hexyl and phenyl magnesium bromide and following the procedure outlined in Example 27.




Table 7 (LCMS² Data: Molecular ion and retention time)

R	R'MgX						
	methyl	ethyl	Isobutyl	4-methoxy phenyl	hexyl	phenyl	n-propyl
methyl			Ex. 27 (326 M+H); 3.50 min	Ex. 28 (376 M+H); 3.13 min	Ex. 29 (354 M+H); 4.05 min		
ethyl		Ex. 30 (312 M+H); 3.18 min	Ex. 31 (340 M+H); 3.69 min	Ex. 32 (390 M+H); 3.32 min			
n-propyl	Ex. 33 (312 M+H); 3.26 min	Ex. 34 (326 M+H); 3.49 min	Ex. 35 (354 M+H); 3.93 min	Ex. 36 (404 M+H); 3.66 min	Ex. 37 (382 M+H); 4.46 min	Ex. 38 (374 M+H); 3.744 min	Ex.39 (340 M+H); 3.6 min
pentyl	Ex. 40 (340 M+H); 3.79 min	Ex. 41 (354 M+H); 4.00 min	Ex. 42 (382 M+H); 4.39 min	Ex. 43 (432 M+H); 4.11 min	Ex. 44 (410 M+H); 4.57 min	Ex. 45 (402 M+H); 4.185 min	
phenyl	Ex. 46 (346 M+H); 3.27 min		Ex. 47 (388 M+H); 3.91 min				
2-furyl	Ex. 48 (336 M+H); 3.00 min	Ex. 49 (350 M+H); 3.28 min	Ex. 50 (378 M+H); 3.69 min		Ex. 51 (406 M+H); 4.19 min		
i-propyl					Ex. 52 (382 M+H); 4.47 min		
biphenyl	Ex. 53 (422 M+H); 3.90 min	Ex. 54 (436 M+H); 4.14 min	Ex. 55 (464 M+H); 4.46 min				

The following compounds (Examples 56-76, Table 8) were prepared using 5-bromothiophene-2-sulfonyl chloride with crotonic acid, 2-pentenoic acid, 2-hexenoic acid, 2-octenoic acid, cinnamic acid, β -(3-pyridyl)-acrylic acid, furylacrylic acid, 4-methyl-2-pentenoic acid, and 4-phenylcinnamic acid and methyl, ethyl, isobutyl, 4-

methoxyphenyl, hexyl and phenyl magnesium bromide and following the procedure outlined in Example 27.






Table 8 (LCMS² Data: Molecular ion and retention time)

R	R'MgX				
	methyl	ethyl	isobutyl	4-methoxy phenyl	Hexyl
methyl			Ex. 56 (372 M+H); 3.52 min		Ex. 57 (400 M+H); 4.07 min
ethyl		Ex. 58 (358 M+H); 3.26 min	Ex. 59 (386 M+H); 3.71 min		
n-propyl		Ex. 60 (372 M+H); 3.52 min	Ex. 61 (400 M+H); 3.95 min		
pentyl		Ex. 62 (400 M+H); 4.02 min	Ex. 63 (428 M+H); 4.41 min	Ex. 64 (478 M+H); 4.12 min	Ex. 65 (456 M+H); 4.57 min
phenyl	Ex. 66 (392 M+H); 3.31 min	Ex. 67 (405 M+H); 3.55 min	Ex. 68 (434 M+H); 3.93 min		
pyridyl			Ex. 69 (433 M+H); 2.67 min		
2-furyl	Ex. 70 (382 M+H); 3.04 min	Ex. 71 (395 M+H); 3.32 min	Ex. 72 (424 M+H); 3.71 min		Ex. 73 (452 M+H); 4.21 min
i-propyl		Ex. 74 (372 M+H); 3.49 min	Ex. 75 (400 M+H); 3.96 min		
biphenyl		Ex. 76 (482 M+H); 4.16 min			

Example 77A**5-Chloro-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide**


Following the procedure outlined in Example 27 (Part 1 and 2), 2-pentenoic
 5 acid was coupled with 4R-4-benzyl-2-oxazolidinone to give R-3-(2'-pentenyl)-4-benzyl-2-oxazolidinone. Addition of hexyl magnesium bromide was followed by trapping by N-bromosuccinimide. After workup, flash chromatography over silica gel using 5% ether in hexane, gave approximately a 2:1 mixture of (1R-2R)- : (1R-2S)-3-(2'-bromo-3'ethylnonanyl)-4-benzyl-2-oxazolidinone.

10 Each isomer was converted to the corresponding sulfonylated amino alcohol following the procedure in Example 27, (Steps 3-5).



15 **Table 9 (LCMS² Data: Molecular ion and retention time)**

	5-chlorothiophene-2-sulfonyl	5-bromothiophene-2-sulfonyl
1S-2R	Example 77A (368 M+H) 4.24 min	Example 78A (414 M+H) 4.26 min
1S-2S	Example 77B (368 M+H) 4.24 min	Example 78B (414 M+H) 4.26 min

Example 79**5-Chloro-N-[(1S)-1-(hydroxymethyl)-2-(methylamino)butyl]thiophene-2-sulfonamide****5 A. Part 1:**

To a solution of D-serine (1.05 g, 10 mmol) in H₂O/THF (1:1, 100 mL) at 0° C was added sodium hydroxide (2.17 g, 30 mmol) and 5-chlorothiophene-2-sulfonyl chloride (2.17 g, 10 mmol). The solution was stirred for 2 to 3 h, then the organic phase was concentrated and the aqueous phase acidified with 1 N HCl and
10 extracted into ethyl acetate and concentrated.

B. Part 2:


To the residue from Part 1 (2.5 mmol) dissolved in THF (25 mL) at -78° C was added ethyl magnesium bromide (7.5 mL, 7.5 mmol). The mixture was warmed to 25° C and allowed to stir for 48 h. It was then acidified with 1 N HCl and
15 extracted into ethyl acetate and concentrated.

C. Part 3:

To the product from Part 2 (0.1 mmol) dissolved in DMF (500 µL) was added CH₂Cl₂ (1.5 mL), acetic acid (12 µL, 0.2 mmol) and methyl amine (2 M solution in THF) (100 µL, 0.2 mmol). The reaction was stirred for 5 min and sodium triacetoxyborohydride (105.6 mg, 0.5 mmol) was added. The solution was allowed to
20 stir for 8-16 h and purified by RP-HPLC¹ to give **Example 79** (6.8 mg).

The following compounds (**Examples 79-86, Table 10**) were prepared using methyl, ethyl or pentyl magnesium bromide with methylamine (2M soln in THF),


ethylamine (2M soln in THF), ethanolamine, benzylamine, and cyclopentylamine and following the procedure outlined in Example 79.



5

Table 10 (LCMS² Data: Molecular ion and retention time)

	R'MgBr		
NHR''	methyl	ethyl	pentyl
Methylamine		Example 79 (313.0 M+H); 1.67 min	
ethylamine	Example 80 (313.0 M+H); 1.53 min		
Ethanolamine	Example 81 (329.0 M+H); 1.22 min	Example 82 (343.0 M+H); 1.73 min	Example 83 (385.0 M+H); 2.36 min
Benzylamine	Example 84 (375.0 M+H); 2.12 min	Example 85 (389.0 M+H); 2.25 min	
cyclopentylamine	Example 86 (353.0 M+H); 1.99 min		

Example 87**5-Chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-N-(2-phenoxyethyl)thiophene-2-sulfonamide****5 A. Part 1**

To a solution of L-isoleucine methyl ester hydrochloride (1.82 g, 10 mmol) and 5-chlorothiophene-2-sulfonyl chloride (1.82 g, 10 mmol) was added triethylamine (4.18 mL, 30 mmol). The mixture was stirred at 60° C overnight, then filtered and concentrated. The crude product was purified by flash chromatography over silica gel using 10% ethyl acetate in hexane to give 5-chlorothiophene-2-sulfonyl isoleucine methyl ester 2.53 g.


B. Part 2

To a solution of 5-chlorothiophene-2-sulfonyl isoleucine methyl ester (103 mg, 0.25 mmol) in DMF (1 mL) was added β-bromophenetole (55 mg, 0.5 mmol) and potassium carbonate (103 mg, 0.75 mmol). The reaction was shaken at 25° C. overnight, then concentrated.

C. Part 3

The residue from part 2 was dissolved in 5% methanol in THF (1 mL) and lithium borohydride (11 mg, 0.5 mmol) added. The reaction was shaken at 25° C. for 2 days then quenched by addition of water (1 mL) and extracted into ethyl acetate (3.5 mL). The organic phase was evaporated and the residue purified by RP-HPLC¹ to give **Example 87** (48 mg).

The following compounds (Examples 87-88, Table 11) were prepared using β -bromophenetole, 3-chlorobenzyl bromide and following the procedure outlined in Example 87.




5

Table 11 (LCMS² Data: Molecular ion and retention time)

	RSO₂Cl
R₁Br	5-chlorothiophene-2-sulfonyl chloride
β -bromophenetole	Example 87 (418.0 M+H); 4.05 min
3-chlorobenzyl bromide	Example 88 (422.0 M+H); 4.12 min

Example 89

5-Chloro-*N*-[(*S*)-2-hydroxy-1-phenylethyl]thiophene-2-sulfonamide



10

To a solution of (*S*)-(+)-2-phenylglycinol (6.8 mg, 0.05 mmol) in CH₃CN (200 μ L) was added Et₃N (105 μ L, 1M in CH₃CN) and 5-chlorothiophene-2-sulfonyl chloride (10.9 mg, 0.05 mmol) as a solution in CH₃CN (200 μ L). The vial was capped

and shaken for 8 to 12 h at 40 °C. Solvent was removed *in vacuo*, and the residue dissolved in 1.6 mL DMSO (0.03 M).

The following compounds (Examples 89-117, Table 12) were prepared using 5-chlorothiophene-2-sulfonyl chloride and 5-bromothiophene-2-sulfonyl chloride with (S)-(+)-2-phenylglycinol, L-leucinol, DL-2-amino-1-hexanol, 2-amino-2-methyl-1-propanol, 2-amino-2-ethyl-1,3-propanediol, cycloleucinol, (S)-cyclohexylalaninol, L-phenylalaninol, L-methioninol, DL-2-amino-1-pentanol, L-*tert*-leucinol, chloramphenicol, (S)-(+)-2-amino-1-butanol, (S)-benzyl-L-cysteinol, benzyl-L-threoninol, 4-methylbenzyl-H-cysteinol, benzyl-H-tyrosinol, and L-threoninol following the procedure outlined in Example 89.




Table 12 (LCMS Data: Molecular ion and retention time)


Amine	X = Cl	X = Br
(S)-(+)-2-phenylglycinol	Example 89 (316.46 M-H), 0.95 min	Example 104 (361.31 M-H); 0.98 min
L-leucinol	Example 90 (296.48 M-H), 1.01 min	Example 105 (342.41 M-H); 1.02 min
DL-2-amino-1-hexanol	Example 91 (296.49 M-H), 1.02 min	Example 106 (342.39 M-H); 1.04 min
2-amino-2-methyl-1-propanol	Example 92 (268.45 M-H), 0.81 min	Example 107 (314.38 M-H); 0.83 min
2-amino-2-ethyl-1,3-propanediol	Example 93 (298.46 M-H), 0.69 min	Example 108 (344.37 M-H); 0.69 min
cycloleucinol	Example 94 (295.02 M-H), 0.92 min	Example 109 (340.4 M-H); 0.93 min
(S)-cyclohexylalaninol	Example 95 (336.31 M-H), 1.19 min	Example 110 (382.41 M-H); 1.2 min

Table 12 (cont'd)

Amine	X = Cl	X = Br
L-phenylalaninol	Example 96 (330.50 M-H), 1.03 min	
L-methioninol		Example 111 (360.33 M-H); 0.9 min
DL-2-amino-1-pentanol	Example 97 (282.68 M-H), 0.92 min	(Example 112 327.07 M-H); 0.94 min
L- <i>tert</i> -leucinol	Example 98 (296.50 M-H), 1.22 min	Example 113 (341.44 M-H); 1.01 min
Chloramphenicol	Example 99 (391.44 M-H), 0.89 min	
(S)-(+)-2-amino-1-butanol	Example 100 (268.45 M-H), 0.61 min	
S-benzyl-L-cysteinol	Example 101 (377.77 M-H), 1.13 min	Example 114 (422.35 M-H); 1.15 min
benzyl-L-threoninol	Example 102 (374.49 M-H), 1.21 min	
4-methylbenzyl-H-cysteinol		Example 115 (436.31 M-H); 1.01 min
benzyl-H-tyrosinol		Example 116 (480.43 M-H); 1.15 min
L-threoninol	Example 103 284.10 (M-H), 0.58 min	Example 117 329.99 (M-H), 0.68 min

Example 118

5-Chloro-N-[(S,S)-1-formyl-2-methylbutyl]thiophene-2-sulfonamide



A. Part 1


To a solution of 5-chlorothiophene-2-sulfonyl chloride (11 g, 50.7 mmol) in CH₃CN (100 mL) and (S)-isoleucinol (6.2 g, 53 mmol) was added Et₃N (11 mL, 109 mmol). The reaction mixture was heated at 50 °C with stirring for 24 h. The solvent was removed and the oil was dissolved in EtOAc (100 mL). The solution was washed with water (2 x 100 mL), brine (1 x 100 mL), and dried over Na₂SO₄. The solvent was removed to give 13.85 g (88%) of the desired sulfonamide.

10 B. Part 2

Molecular sieves (15 g, 4 Å) were stirred in dry CH₂Cl₂ (175 mL) for 10 min. Then a mixture of pyridinium chlorochromate (8.6 g, 39.9 mmol) and silica gel (9 g) was added and the mixture was stirred an additional 10 min. To the suspension was added 5-chlorothiophene-2-sulfonyl isoleucinol (4 g, 13.4 mmol) dissolved in CH₂Cl₂ (15 mL) and the resulting slurry was stirred for 2 h. The reaction mixture was filtered and the solvent was removed. The residue was subjected to a Biotage™ eluting with 20% EtOAc/hexane to give 3.22 g (81%) of the aldehyde (LCMS = 294.21 (M-H), rt = 1.10 min).

Example 119


5-Chloro-N-[(S,S)-1-(1-hydroxyethyl)-2-methylbutyl]thiophene-2-sulfonamide



5 To a solution of the aldehyde from example 118 (23.7 mg, 0.08 mmol) in THF (400 μ L) was added methyl magnesium bromide (400 μ L, 1.0 M in THF, 5 eq). The vial was capped and agitated at 50 $^{\circ}$ C for 12h. The reaction was quenched with sat. aqueous NH_4Cl (1.5 mL) and EtOAc (1 mL). The organic layer was transferred into a tarred vial and the aqueous layer was extracted with EtOAc (1 mL). The combined
10 organics were concentrated (Savant, medium heat) and the resulting mixture of diastereomers was dissolved in DMSO such that the final concentration was 30 mM.

The following compounds (Examples 119-154, Table 13) were prepared using 5-chlorothiophene-2-sulfonyl isoleucinal (example 118) and 5-bromothiophene-2-sulfonyl isoleucinal (prepared as in example 118) with methylmagnesium bromide,
15 cyclopentylmagnesium bromide, hexylmagnesium bromide, pentylmagnesium bromide, butylmagnesium bromide, isopropylmagnesium bromide, *o*-tolylmagnesium bromide, *tert*-butylmagnesium bromide, isobutylmagnesium bromide, vinylmagnesium bromide, allylmagnesium bromide, ethylmagnesium bromide, 4-fluorophenylmagnesium bromide, 4-chlorophenylmagnesium bromide, 2-methyl-1-
20 propenylmagnesium bromide, isopropenylmagnesium bromide, 4-anisylmagnesium bromide, 1-methyl-1-propenylmagnesium bromide, 2-[2-(1,3-dioxanyl)]ethylmagnesium bromide, 3-butenylmagnesium bromide, 1-propynylmagnesium bromide, 4-thioanisolemagnesium bromide, and 4-*N,N*-dimethylanilinemagnesium bromide following the procedure outlined in example 119.

Note: during the reaction sequence with the 5-bromothiophene compounds, the bromine is converted to a hydrogen on the thiophene ring.



5


Table 13 (LCMS Data: Molecular ion and retention time)

R-MgBr	5-chlorothiophene	thiophene
methylmagnesium bromide	Ex. 119	
	310.09(M-H), 1.06 min	
	310.10 (M-H), 1.12 min	
cyclopentylmagnesium bromide	Ex. 120	Ex. 140
	364.13 (M-H), 1.41 min	330.19 (M-H), 1.26 min
Hexylmagnesium bromide	Ex. 121	Ex. 141
	380.16 (M-H), 1.50 min	346.24 (M-H), 1.38 min
	380.17 (M-H), 1.54 min	346.24 (M-H), 1.42 min
pentylmagnesium bromide	Ex. 122	Ex. 142
	366.15 (M-H), 1.42 min	332.19 (M-H), 1.30 min
	366.16 (M-H), 1.47 min	332.19 (M-H), 1.35 min
Butylmagnesium bromide	Ex. 123	Ex. 143
	352.15 (M-H), 1.34 min	318.18 (M-H), 1.26 min
	352.13 (M-H), 1.40 min	
isopropylmagnesium bromide	Ex. 124	
	338.11 (M-H), 1.31 min	
o-tolylmagnesium bromide		Ex. 144
		352.16 (M-H), 1.24 min
tert-butylmagnesium bromide	Ex. 125	Ex. 145
	352.14 (M-H), 1.41 min	318.2 (M-H), 1.28 min
isobutylmagnesium bromide	Ex. 126	
	352.14 (M-H), 1.33 min	
	352.13 (M-H), 1.38 min	
vinylmagnesium bromide	Ex. 127	Ex. 146
	322.09 (M-H), 1.14 min	288.15 (M-H), 0.98 min
	322.10 (M-H), 1.19 min	288.15 (M-H), 1.02 min
allylmagnesium bromide	Ex. 128	Ex. 147
	336.11 (M-H), 1.22 min	302.17 (M-H), 1.06 min
	336.12 (M-H), 1.27 min	302.17 (M-H), 1.11 min
ethylmagnesium bromide	Ex. 129	Ex. 148
	324.10 (M-H), 1.18 min	290.18 (M-H), 1.01 min
	324.11 (M-H), 1.22 min	290.17 (M-H), 1.06 min

Table 13 (cont'd)

R-MgBr	5-chlorothiophene	Thiophene
4-chlorophenylmagnesium bromide	Ex. 131 406.06 (M-H), 1.36 min 406.06 (M-H), 1.41 min	
2-methyl-1-propenylmagnesium bromide	Ex. 132 350.13 (M-H), 1.25 min 350.10 (M-H), 1.31 min	
isopropenylmagnesium bromide	Ex. 133 336.11 (M-H), 1.25 min 336.10 (M-H), 1.28 min	
4-anisylmagnesium bromide	Ex. 134 402.13 (M-H), 1.25 min 402.12 (M-H), 1.31 min	Ex. 149 368.16 (M-H), 1.14 min 368.16 (M-H), 1.16 min
1-methyl-1-propenylmagnesium bromide	Ex. 135 350.13 (M-H), 1.27 min 350.12 (M-H), 1.35 min	
2-[2-(1,3-dioxanyl)]ethylmagnesium bromide	Ex. 136 410.15 (M-H), 1.15 min	Ex. 150 376.19 (M-H), 0.98 min
3-butenylmagnesium bromide	Ex. 137 350.11 (M-H), 1.30 min 350.12 (M-H), 1.33 min	Ex. 151 316.17 (M-H), 1.15 min 316.17 (M-H), 1.18 min
1-propynylmagnesium bromide		Ex. 152 300.17 (M-H), 1.20 min (M-H), min
4-thioanisolemagnesium bromide	Ex. 138 418.11 (M-H), 1.35 min 418.11 (M-H), 1.39 min	Ex. 153 384.13 (M-H), 1.26 min (M-H), min
4- <i>N,N</i> -dimethylanilinemagnesium bromide	Ex. 139 415.15 (M-H), 0.89 min 415.17 (M-H), 0.93 min	Ex. 154 381.21 (M-H), 0.68 min 381.21 (M-H), 0.71 min

Example 155

5-Chloro-*N*-{[(*S,S*)-1-[(*S*)-cyclohex-2-en-1-yl(hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide

To a two dram vial containing magnesium turnings (60 mg, 2.5 mmol)
 5 suspended in THF (3 mL) was added 2-bromocyclohexene (288 μ L, 2.5 mmol)
 followed by 5-chlorothiophene-2-sulfonyl isoleucinal (1 mL of a 1 M THF solution, 1
 mmol, example 118). The vial was capped and agitated at 50 $^{\circ}$ C for 18 h. The vial was
 cooled and sat. aqueous NH_4Cl (1 mL) was added. The vial was vortexed and the
 organic layer was transferred into a tarred vial and the aqueous layer was extracted
 10 with EtOAc (1 mL). The combined organics were concentrated *in vacuo* and the
 residue was submitted to semipreparative RP-HPLC using the conditions below.

Semi-preparative RP-HPLC conditions.

Column: Spring Axial compression; Kromasil C18 10 μ m particle size; 50x150 mm

15 Solvent A: Water (0.1% TFA)

Solvent B: Acetonitrile

Solvent Gradient: 15-95% over 24 min, full cycle is 35 min

Flow Rate: 60ml/min

The product peak was collected based on UV (or ELSD) absorption.

20

The following compounds (Examples 155-161, Table 14) were prepared
 using 2-bromocyclohexene, crotyl bromide, 1-bromo-2-pentene, 3-bromo-2-
 methylpropene, and cinnamyl bromide following the procedure outlined in Example
 155.

25





Table 14 (LCMS Data: Molecular ion and retention time)

Alkyl bromide	
2-bromocyclohexene	Example 155 376.70 (M-H), 1.27 min
crotyl bromide	Example 156 350.50 (M-H), 1.22 min
crotyl bromide	Example 157 350.70 (M-H), 1.25 min
1-bromo-2-pentene	Example 158 364.60 (M-H), 1.35 min
3-bromo-2-methylpropene	Example 159 350.60 (M-H), 1.20 min
3-bromo-2-methylpropene	Example 160 350.60 (M-H), 1.24 min
cinnamyl bromide	Example 161 412.60 (M-H), 1.34min

Example 162**5-Chloro-N-[(S,S)-1-(1-hydroxy-1-methylethyl)-2-methylbutyl]thiophene-2-sulfonamide****5 A. Part 1**

To a solution of 5-chlorothiophene-2-sulfonyl chloride (1.09 g, 5 mmol) in CH₃CN (20 mL) was added (L)-isoleucine methyl ester hydrochloride (908.5 mg, 5 mmol) as a solution in CH₃CN (10 mL) and Et₃N (1 mL, 7.2 mmol). The reaction mixture was heated at 50 °C with shaking for 3 days. The solvent was removed and the oil was dissolved in EtOAc (10 mL). The solution was washed with water (5 mL), sat. NH₄OH (5 mL), and brine (5 mL), and dried over MgSO₄. The solvent was removed to give 1.44 g (88%) of the desired sulfonamide.

B. Part 2

To a solution of the ester from part 1 (40.7 mg, 0.125 mmol) in THF (500 µL) was added methyl magnesium bromide (333 µL, 3.0 M in THF, 8 eq). The vial was capped and agitated at 50 °C for 12h. The reaction was quenched with sat. aqueous NH₄Cl (1.5 mL) and EtOAc (1 mL). The organic layer was transferred into a tarred vial and the aqueous layer was extracted with EtOAc (1 mL). The combined organics were concentrated (Savant, medium heat) and the product was dissolved in DMSO such that the final concentration was 30 mM.

The following compounds (**Examples 162-176, Table 15**) were prepared using 5-chlorothiophene-2-sulfonyl isoleucine methyl ester and 5-bromothiophene-2-sulfonyl isoleucine methyl ester (from part 2) with methylmagnesium bromide, pentylmagnesium bromide, phenylmagnesium bromide, allylmagnesium bromide, ethylmagnesium bromide, 4-chlorophenylmagnesium

bromide, isopropenylmagnesium bromide, 4-anisylmagnesium bromide, 1-methyl-1-propenylmagnesium bromide, 3-butenylmagnesium bromide, 1-propynylmagnesium bromide, 1-naphthylmagnesium bromide following the procedure outlined in example 162.

5





Table 15 (LCMS Data: Molecular ion and retention time)

R-MgBr	X = Cl	X = Br
methylmagnesium bromide	Example 162 324.58 (M-H), 1.19 min	
pentylmagnesium bromide	Example 163 436.69 (M-H), 1.79 min	
phenylmagnesium bromide	Example 164 448.60 (M-H), 1.50 min	
Allylmagnesium bromide	Example 165 376.58 (M-H), 1.44 min	
ethylmagnesium bromide	Example 166 352.63 (M-H), 1.39 min	Example 173 395.49 (M-H), 1.45 min
4-chlorophenylmagnesium bromide	Example 167 516.49 (M-H), 1.62 min	
isopropenylmagnesium bromide	Example 168 376.60 (M-H), 1.51 min	Example 174 421.29 (M-H), 1.39 min
4-anisylmagnesium bromide	Example 169 508.59 (M-H), 1.45 min	
1-methyl-1-propenylmagnesium bromide	Example 170 404.61 (M-H), 1.65 min	Example 175 447.46 (M-H), 1.52 min
3-butenylmagnesium bromide	Example 171 404.64 (M-H), 1.54 min	Example 176 447.25 (M-H), 1.26 min
1-naphthylmagnesium bromide	Example 172 548.56 (M-H), 1.64 min	

Example 177**5-Chloro-N-[1-(hydroxymethyl)cyclohexyl]thiophene-2-sulfonamide****A. Part 1**

To a suspension of 1-amino-1-cyclohexane carboxylic acid (5 g, 35
 5 mmol) and THF (100 mL) was added borane dimethyl sulfide (50 mL, 2M in THF) at
 0 °C. The cold bath was allowed to expire and the reaction was stirred at 25°C
 overnight. NaOH (3M, 100 mL) was added and the mixture was stirred for 4 h. The
 reaction mixture was saturated with K₂CO₃ and extracted with Et₂O (2 x 100 mL).
 The combined organics were washed with brine (100 mL) and dried over MgSO₄ to
 10 give 4.35 g (96%) of the desired amino alcohol.

B. Part 2

The amino alcohol was sulfonylated as in Example 89.

The following compounds (**Examples 177-183, Table 16**) were prepared
 15 using the amino alcohols of 1-amino-1-cyclohexane carboxylic acid, 2-amino-2-
 norbornane carboxylic acid, D,L-1-aminoindane-1-carboxylic acid, and 2-
 aminoindane-2-carboxylic acid hydrochloride with 5-chlorothiophene-2-sulfonyl
 chloride and 5-bromothiophene-2-sulfonyl chloride following the procedure outlined
 for Example 177.

20




Table 16 (LCMS Data: Molecular ion and retention time)

Amino acid	X = Cl	X = Br
1-amino-1-cyclohexane carboxylic acid	Example 177 308.14 (M-H), 1.00 min	Example 181 353.99 (M-H), 1.02 min
2-amino-2-norbornane carboxylic acid	Example 178 320.13 (M-H), 1.04 min	Example 182 366.03 (M-H), 1.06 min
D,L-1-aminoindane-1-carboxylic acid	Example 179 342.12 (M-H), 1.09 min	
2-aminoindane-2-carboxylic acid HCl	Example 180 342.12 (M-H), 1.07 min	Example 183 388.01 (M-H), 1.08 min

The following compounds (Examples 184-195, Table 17) were synthesized using 5-chlorothiophene-2-sulfonyl isoleucinal and 5-bromothiophene-2-sulfonyl isoleucinal with methylmagnesium bromide, *n*-propylmagnesium chloride, and allylmagnesium bromide following the procedure outlined for example 119. The resulting mixtures of diastereomers were isolated by semi-preparative RP-HPLC using the conditions outlined for example 155.

10

Table 17 (LCMS Data³: Molecular ion and retention time)

R-MgX	X = Cl	X = Br
methylmagnesium bromide	Example 184 310.41 (M-H), 0.87 min	Example 190 356.20 (M-H), 0.95 min
	Example 185 310.44 (M-H), 0.93 min	Example 191 356.15 (M-H), 1.02 min
<i>n</i> -propylmagnesium chloride	Example 186 338.47 (M-H), 1.06 min	Example 192 384.17 (M-H), 1.13 min
	Example 187 338.45 (M-H), 1.12 min	Example 193 384.24 (M-H), 1.18 min
allylmagnesium bromide	Example 188 336.10 (M-H), 1.2 min	Example 194 382.00 (M-H), 1.2 min
	Example 189 336.10 (M-H), 1.24 min	Example 195 382.00 (M-H), 1.25 min

The pure synthetic diastereomer of Example 189 was prepared as follows.

A. Part 1

To a -78°C solution of the Weinreb amide (see: F. Roux, et. Al. *Tetrahedron*, 1994, 50 (18), 5345-5360) of BOC-protected isoleucine (13.17 g, 48 mmol) was added allylmagnesium bromide (90 mL, 1M in THF). The cold bath was allowed to expire and the reaction stirred at 25°C overnight. The reaction was quenched by the addition of cold aq. HCl (150 mL, 1M). After 30 min of stirring, the layers were separated and the aqueous layer extracted with ethyl acetate (3 x 75 mL). The combined organics were dried over MgSO_4 and the solvent removed to give 8.41 g (69 %) of the desired ketone.

B. Part 2

To a solution of the ketone from part 1 (8.4 g) in MeOH (200 mL) was added NaBH_4 (1.5 g, 39.6 mmol) as a solid. The reaction was stirred at 25°C for 5 h at which time the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate (100 mL) and washed with water (2 x 50 mL). The crude product was subjected to a Biotage™ eluting with 5 to 15 % EtOAc/hexane to give 4.12 g (49 %) of the desired alcohol.

C. Part 3

A solution of alcohol from part 2 (4.12 g, 16 mmol), CH_2Cl_2 (75 mL), and TFA (15 mL) was stirred at 25°C for 15 min. The reaction was quenched with a solution of NaOH (15 mL, 1M) and then basified to pH 12 with NaOH pellets. The resulting solution was extracted with CH_2Cl_2 (2 x 50 mL), and the combined organics were washed with water (25 mL), brine (25 mL), and dried over MgSO_4 to give 2.44 g (97 %) of the desired amino alcohol, which was carried on to part 4 without further purification.

D. Part 4

The amino alcohol was sulfonylated as in example 89. The pure synthetic diastereomer of Example 193 was obtained as follows:


To a solution of the BOC-amino homoallyl alcohol (1.1 g, 4.27 mmol, see part 1-2 of example 188) in absolute EtOH (50 mL) was added Pd/C (110 mg). The flask was placed under an atmosphere of hydrogen (balloon) and stirred at 25°C . Following completion of the reaction (2h), the mixture was filtered through a pad of

Celite and the solvent was removed to give 1.16 g (quant) of the propyl analog. The BOC group was removed and the amine sulfonylated per the protocol outlined for example 189.

5


Example 196

5-Chloro-*N*-[(*S,S*)-2-methyl-1-(2,2,2-trifluoro-1-hydroxyethyl)butyl]thiophene-2-sulfonamide



To a 0 °C solution of 5-chlorothiophene-2-sulfonyl isoleucinal (770 mg, 2.6 mmol, see Example 118, parts 1&2) in THF (5 mL) was added TMS-CF₃ (5 mL, 0.5M in THF). The resulting mixture was treated with TBAF (250 µL, 1M in THF). The cold bath was removed and the reaction was stirred at 25°C overnight. The reaction was quenched with HCl (25 mL, 2M) and the resulting solution was extracted with ethyl acetate (3 x 15 mL). The combined organic extracts were washed with water (25 mL) and brine (25 mL) then dried over MgSO₄. The residue was submitted to RP-HPLC (see example 155 for procedure) to give 74 mg of the desired product (m/z = 364.0 (M-H), rt = 1.23 min).

The following compounds (Examples 197-198, Table 18) were synthesized using 1-amino-1-cyclohexane carboxylic acid and 5-chlorothiophene-2-sulfonyl chloride with allylmagnesium bromide and 2-methyl allylmagnesium chloride following the four step procedure outlined for examples 177 (parts 1&2), 118 (part 2), and 119 respectively.


**Table 18 (LCMS Data: Molecular ion and retention time)**

R-MgX	
allylmagnesium bromide	Example 197 348.10 (M-H), 1.18 min
2-methylallylmagnesium chloride	Example 198 364.10 (M-H), 1.26 min

5

Example 199A

5-Chloro-N-[(S)-2-hydroxy-1-(4-hydroxycyclohexyl)ethyl]thiophene-2-sulfonamide




mmoles). The reaction was allowed to warm up to 25°C and stirred overnight. The dioxane and Et₃N were removed and the remaining aqueous solution diluted with 1N aq. HCl. The resulting precipitate was collected, washed with water and diethyl ether to give the desired product as a white solid (12g, 50% in two steps) (100% purity by ELSD, m/z= 352 (M-1)).

B. Part 2

To a suspension of (S)-N-(5-Chlorothiophene-2-sulfonyl)-4-hydroxycyclohexylglycine (12 g, 33.99 mmol, part 1) in anhydrous THF was added borane-THF (110 mL, 1 M in THF, 110 mmoles) dropwise at 0°C. The resulting mixture was stirred at 25°C over the weekend. The reaction mixture was quenched with HCl (75 mL, 1M) at 0 °C and stirred at 25°C for 1 h. THF was removed and the precipitate was collected, washed with water (containing a small amount of diethyl ether), and dried to give a white solid as the desired product (9 g, 78%) (100% purity by ELSD, m/z= 338.5 (M-1), HPLC retention time³=0.64 min).

Example 199

5-Chloro-N-[(S)-2-hydroxy-1-(4-methoxycyclohexyl)ethyl]thiophene-2-sulfonamide



A. Part 1

A mixture of 5-chloro-N-[(S)-2-hydroxy-1-(4-hydroxycyclohexyl)ethyl]thiophene-2-sulfonamide (6.4 g, 18.83 mmol) from Example 199A, 2,2-dimethoxypropane (7 mL, 5.65 mmol), and TsOH·H₂O (72 mg,

0.38 mmol.) in anhydrous benzene (120 mL) was refluxed. After one hour, benzene was slowly distilled under atmospheric pressure to a final volume of 10 mL. Fresh benzene (100 mL) and 2,2-dimethoxypropane (5 mL) were added and the above operation was repeated. The residue was partitioned between diethyl ether and sat.

- 5 NaHCO_3 . The aqueous layer was extracted with diethyl ether (3 x 100 mL), and the combined extracts were dried over MgSO_4 . The crude product was purified by column chromatography using 1:5 EtOAc/ CH_2Cl_2 as an eluent to give the *N,O*-acetonide (5.77 g, 81%) (100%, $m/z=380$ ($M+1$)).

B. Part 2

- 10 To a 0°C solution of (*S*)-2-(5-chlorothiophene-2-sulfonamido)-2-(4-hydroxycyclohexyl)-*N,O*-acetonide (379 mg, 1 mmol) in THF (7 mL) and DMF (2 mL) was added NaH (80 mg, 2 mmol). The resulting reaction was stirred at 0°C for 10 min at which time iodomethane (311 μL , 5 mmol) was added. The reaction was allowed to warm to 25°C and stirred for 18 h. The solvent was removed and acetic acid (80%, 15 mL) was added. The mixture was stirred at 25°C over the weekend. Following removal of the acetic acid *in vacuo*, the residue was subjected to column chromatography on silica gel eluting with MeOH/ CH_2Cl_2 (3:10) to give 303 mg (86 %) of the desired product.

- 20 The following compounds (Examples 199-202B, Table 19) were prepared using (*S*)-2-(5-chlorothiophenesulfonamido)-2-(4-hydroxycyclohexyl)-*N,O*-acetonide (from Example 199, part 1) with iodomethane, 1-bromopropane, allyl bromide, benzyl bromide, 2-picolyl chloride hydrochloride and 3-picolyl chloride hydrochloride as outlined in Example 199.





Table 19 (LCMS Data³: Molecular ion and retention time)

R-X	
iodomethane	Example 199 352.1(M-H), 0.82min
1-bromopropane	Example 200 366.0(M-H), 0.91min
allyl bromide	Example 201 378.0(M-H), 1.01min
benzyl bromide	Example 202 428.1(M-H), 1.21min
2-picolyl chloride HCl	Example 202A 429.4(M-H), 0.57min
3-picolyl chloride HCl	Example 202B 429.0(M-H), 0.57min

Example 203

5 N-[1-Acetyl-4-(hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide.



A. N-[1-Boc-4-(Carboxylic acid)piperidin-4-yl]-5-chlorothiophene-sulfonamide

- 10 Triethylamine (2.28 mL, 1.66g, 16.45 mmol) was added to a slurry of 1-Boc-4-aminopiperidine-4-carboxylic acid (2.68g, 10.973 mmole) in acetonitrile:water (1:1) (40 mL) at 25°C. The slurry became a neon yellow to greenish solution at the end of addition. The slurry was slightly warmed up (5 min) in order to obtain a solution. The mixture was cooled to 0°C, 5-chlorothiophene-2-sulphonyl chloride (2.62g, 12.07 mmol) was added (5 min) dropwise as a solution in acetonitrile (8 mL). The solution was allowed to warm up to 25°C overnight. After 19 h, an aliquot was taken. TLC (9:1 CH₂Cl₂:CH₃OH) indicated that reaction was about 90%

done. The reaction was quenched by addition of water (50 mL), CH₂Cl₂ (50 mL) and ice cold 1N HCl (10 mL). The organic layer was washed with water and saturated NaCl. It was dried over MgSO₄, filtered, and concentrated to a yellow oil (2.1g). The crude material was purified by column chromatography, silica gel 230 to 400 mesh, eluent: starting with 5% MeOH in CH₂Cl₂ and ending with 10% MeOH in CH₂Cl₂ to furnish N-[1-Boc-4-(carboxylic acid)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide as a white amorphous solid (1.2g, 25.7%). Mass Spectrum (-ESI): 423 (M-H)⁺.

B. N-[1-Boc-4-(hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide.

10

1N Borane-THF (1.019g, 12.14 mL, 11.86 mmol) was added dropwise over 30 min at 0° C to a solution of N-[1-Boc-4-(carboxylic acid)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide (1.2g, 2.82 mmol) in anhydrous tetrahydrofuran (15 mL). The reaction was allowed to warm up to 25°C overnight, and then was quenched by addition of 30 mL of 10% acetic acid in methanol. After solvent evaporation, the crude product was dissolved in ethyl acetate and washed with 1M HCl, water and 10% NaHCO₃. The organic layer was dried over MgSO₄, filtered and concentrated to obtain a crude yellow oil (1.1g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluent: starting with 1:3 EtOAc-hexane and ending with 1:1 EtOAc-hexane to afford N-[1-Boc-4-(hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide as a colorless oil (0.79g, 68.2%). Mass Spectrum (-ESI): 409 (M-H)⁺.

25

C. N-[4-(Hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide HCl Salt

30


4N HCl (5mL) was added to a stirred solution of N-[1-Boc-4-(hydroxymethyl) piperidin-4-yl]-5-chlorothiophene-2-sulfonamide **5** (0.7g, 1.7 mmol) in EtOAc (4 mL). The solution was allowed to stir at 25°C. After 30 min, a cloudy solution formed. After 2 h a precipitate formed. TLC (1:1 EtOAc-hexane) indicated the reaction was complete. The solvent was reduced to ~2-3 mL, diluted with diethyl ether (6 mL) and filtered through a filter funnel. The precipitate was washed with diethyl ether (3 x 5 mL) to obtain N-[4-(hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide as an amorphous white solid (0.48 g, 90.7%). Mass Spectrum (+ESI): 311 (M+H)⁺.

D. N-[1-Acetyl-4-(hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide

- Acetyl chloride (0.15g, 1.894 mmol) was added dropwise (5 min) as a solution in CH₂Cl₂ (1 mL) to a cold 0° C solution of N-[4-(hydroxymethyl)piperidin-4-yl]-5-chloro- thiophene-2-sulfonamide (0.19g, 0.61 mmol) in CH₂Cl₂ (5 mL) and triethylamine (0.44 mL, 3.18 mmol) The solution was allowed to warm to 25° C overnight (19 h). An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was complete. It was diluted with CH₂Cl₂ (10 mL) and the organic layer was washed with 1N HCl (50 mL), saturated aqueous NaHCO₃ (50 mL) and NaCl (50 mL). The organic layer was dried over MgSO₄, filtered, and concentrated to obtain a crude oil (175 mg). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluent: starting with 1:4 EtOAc-hexane and ending with 1:1 EtOAc-hexane to afford N-[1-Acetyl-4-(hydroxymethyl)-piperidin-4-yl]-5-chlorothiophene-2-sulfonamide as yellowish color oil (62 mg, 28.9%).
- Mass Spectrum (+ESI): 353 (M+H)⁺. Anal. Calc'd for C₁₂H₁₇ClN₂O₄S₂ .1.62 H₂O: C, 37.29; H, 5.70; N, 7.26. Found: C, 37.62; H, 5.36; N, 7.31.

Example 204

5-Chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide.



20

A. 2-Chlorofuran.

1.6 M nBuLi (15.37 g, 150 mL, 0.24 mol) was added dropwise over 10 min at 25° C to a solution of furan (13.6 g, 0.20 mol) in dry diethyl ether (200 mL). When the dropwise addition was completed, the reaction mixture was cooled to

-70°C. At this temperature a solution of hexachloroethane (49.8g, 0.21 mol) was added over 10 min and the temperature was not allowed to rise above -55°C. The reaction mixture was kept at -70°C for 3 h. The reaction mixture was then warmed to 25°C, hydrolyzed with ice water and neutralized with 2.5 N hydrochloric acid. The phases were separated and the water phase extracted twice with diethyl ether (100 mL). The combined diethyl ether phases were washed once with a solution of NaHCO₃ (50 mL) and once with water (50 mL) and dried over MgSO₄. The diethyl ether was distilled off through a fractionating column and the product was collected at 78 to 79°C to obtain 2-chlorofuran as a colorless oil (20.0 g, 97.6 %). ¹H-NMR (DMSO-d₆, 400MHz) δ 7.34 (d, 1H); 6.38 (d, 1H); 6.21 (d, 1H).

B. 5-Chlorofuran-2-sulfonyl chloride

Phosphorus pentachloride (40.53g, 0.1947 mol) was added portionwise (caution, foaming) over 5 min at 25°C to chlorosulfonic acid (56.8g, 32.4 mL, 0.487 mol) and the resulting solution was stirred at 25°C for 10 min. Then, 2-chlorofuran (20.0 g, 0.1947 mol) was added in one portion and the resulting dark suspension was heated to 55°C for 1.0 h during which time foaming occurred and subsided. The reaction mixture was then poured onto ice and the resulting suspension was extracted with CH₂Cl₂ (250 mL). The organic was filtered through a pad of celite, washed with brine (70 mL) and dried over MgSO₄. The solvent was removed *in vacuo* to provide 5-chlorofuran-2-sulfonyl chloride as a black oil (14.1g, 36.02 %). ¹H-NMR (DMSO-d₆, 400MHz) δ 7.05 (d, 1H); 6.35 (d, 1H).

C. 5-Chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide.


5-Chlorofuran-2-sulfonyl chloride (3.376g, 16.79 mmol) was added dropwise (5 min) as a solution in CH₂Cl₂ (10 mL) to a 0°C solution of L-isoleucinol (1.5g, 12.92 mmol) in CH₂Cl₂ (15 mL) and triethylamine (2.69 mL, 19.38 mmol). The solution was allowed to warm to 25°C overnight (19 h). An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was complete. It was diluted with CH₂Cl₂ (100 mL) and the organic layer was washed with 1N HCl (2x50 mL), saturated aqueous NaCl (50 mL). The organic layer was dried over MgSO₄, filtered, and concentrated to obtain a crude black oil (2.69 g). The crude product was purified by column chromatography, silica gel 230 to 400 mesh, eluent starting with 1:4

EtOAc-hexane and ending with 1:1 EtOAc-hexane to afford 5-chloro-N-[(1S,2S-1-(hydroxymethyl)-2-methylbutyl)-2-furansulfonamide as an amorphous white solid (0.98 g, 26.92 %). Mass Spectrum (-ESI): 280 (M-H)⁺. Anal. Calc'd for C₁₀H₁₆ClN₂O₄S: C, 42.63; H, 5.72; N, 4.97. Found: C, 42.34; H, 5.65; N, 4.77.

5

Example 205

N-[(1*S*)-2-Butyl-1-(hydroxymethyl)hexyl]-5-chloro-2-thiophenesulfonamide .



- 10 A. (4*R*)-4-Benzyl-3-[(*E*)-2-heptenoyl]-1,3-oxazolidin-2-one
- Triethylamine (6.85g, 49.15 mmol) and trimethyl acetyl chloride (6.05 mL, 49.15 mmol) were added dropwise (5 min) to a -78°C solution of 2-heptenoic acid (6 g, 46.81 mmole) in THF (80 mL). The slurry was stirred at -78°C for 5 min and then replaced with a 0°C cooling system. It was stirred at this temperature for 1 h.
- 15 In a separate flask, a solution of R-(+)-4-benzyl-2-oxazolidinone (8.295 g, 46.81 mmol) was cooled to -78°C and nBuLi (1.6M, 46.8 mmol) was added dropwise over 10 min. The colorless solution was stirred at this temperature for 45 min and transferred via cannula to a -78°C solution of the ester. The yellowish slurry was warmed to 25°C overnight (19 h). An aliquot was taken and TLC (1:1 EtOAc-hexane)
- 20 indicated that reaction was complete. It was cooled to 0°C and the reaction was quenched by addition of H₂O (20 mL). It was diluted with ethyl acetate (200 mL) and the organic layer was separated. The organic layer was dried over MgSO₄, filtered, and concentrated to obtain a crude yellow oil (13.69 g). The crude product was purified by column chromatography, silica gel 230 to 400 mesh, eluent: 1:4 EtOAc-

hexane to obtain (4*R*)-4-benzyl-3-[(*E*)-2-heptenoyl]-1,3-oxazolidin-2-one as a colorless oil (12.1 g, 92.80 %). Mass Spectrum (-ESI): 288 (M-H)⁺.

5 B. (4*R*)-4-Benzyl-3-[(2*R*)-2-bromo-3-butylheptanoyl]-1,3-oxazolidin-2-one

A slurry of copper bromide (I) dimethyl sulfide complex (5.132 g, 24.967 mmol) in THF (60 mL) and dimethyl sulfide (30 mL) as a co-solvent was cooled to -40°C and *n*-butyl magnesium chloride (25 mL, 49.93 mmol) was added dropwise for 10 min and stirred for 20 min while warming to -15°C. The black slurry was cooled to -40°C and (4*R*)-4-benzyl-3-[(*E*)-2-heptenoyl]-1,3-oxazolidin-2-one (6 g, 20.80 mmol) was added dropwise over 10 min as a solution in THF (20 mL) at -40°C. The reaction was let warm up to 25°C overnight (20 h). *N*-Bromosuccinimide (7.407 g, 41.61 mmol) was added portionwise to a cold -78°C solution of the black slurry. It was allowed to warm to 0°C and was stirred for an additional 3 h. The reaction was quenched with a 1:1 solution of saturated ammonium carbonate and 0.5 N potassium bisulfate. The black slurry became greenish to blue. A precipitate formed (light blue). It was filtered. The mother liquor was diluted with ethyl acetate (150 mL) and the organic was dried over MgSO₄, filtered, concentrated to obtain (4*R*)-4-benzyl-3-[(2*R*)-2-bromo-3-butylheptanoyl]-1,3-oxazolidin-2-one as a crude semi-solid (green) (8.49 g, 96.15 %). Mass Spectrum (-ESI): 423 (M-H)⁺.

 C. (4*R*)-3-[(2*S*)-2-Azido-3-butylheptanoyl]-4-benzyl-1,3-oxazolidin-2-one

Tetramethylguanidine azide (TMGA) (5.398 g, 37.70 mmol) was added dropwise (5 min) to a 25° C solution of (4*R*)-4-benzyl-3-[(2*R*)-2-bromo-3-butylheptanoyl]-1,3-oxazolidin-2-one (4.0 g, 9.42 mmol) in acetonitrile (50 mL). The reaction was stirred for 4 days. An aliquot was taken and TLC (1:4 EtOAc-hexane) indicated that reaction was complete. The solvent was removed *in vacuo*. The resulting black semi-solid was dissolved in CH₂Cl₂ (200 mL) and quenched with 1N HCl (30 mL). The organic layer was dried over MgSO₄, filtered, and concentrated *in vacuo* to obtain (4*R*)-3-[(2*S*)-2-azido-3-butylheptanoyl]-4-benzyl-1,3-oxazolidin-2-one as a crude yellow oil (3.61 g, 99.1 %). Mass Spectrum (-ESI): 385 (M-H)⁺.

D. (2S)-2-Amino-3-butyl-1-heptanol.


To a slurry of LAH (1.219g, 32.13 mmol) in THF (60 mL) was added (4R)-3-[(2S)-2-azido-3-butylheptanoyl]-4-benzyl-1,3-oxazolidin-2-one (3.6g, 9.37 mmol) dropwise at 0°C over 20 min. The reaction was heated to 36°C for 18 h. The reaction slurry (brown) was cooled to 0°C and the reaction was quenched with H₂O (15 mL) and washed with 1N NaOH (30 mL) and H₂O (15 mL). It was let stir for 2 h to obtain an off-white slurry. The slurry was filtered and the mother liquor was further dried over MgSO₄, filtered, and concentrated *in vacuo* to obtain (2S)-2-amino-3-butyl-1-heptanol as a crude yellow oil (1.93g, 73.75%). Mass Spectrum (+ESI): 188 (M+H)⁺.

E. N-[(1S)-2-Butyl-1-(hydroxymethyl)hexyl]-5-chloro-2-thiophenesulfonamide.

5-Chlorothiophene-2-sulfonyl chloride (2.42g, 11.55 mmol) was added dropwise (5min) as a solution in CH₂Cl₂ (20 mL) to a 0°C solution of (4R)-3-[(2S)-2-azido-3-butylheptanoyl]-4-benzyl-1,3-oxazolidin-one (1.9 g, 10.14 mmol) and triethylamine (2.11 mL, 15.21 mmol). The solution was allowed to warm to 25°C overnight (19 h). An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was complete. It was diluted with CH₂Cl₂ (100 mL) and the organic layer was washed with 1N HCl (2x50 mL) and saturated aqueous NaCl (50 mL). The organic layer was dried over MgSO₄, filtered, and concentrated to obtain a crude oil (2.98 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluent: starting with 1:3 EtOAc-hexane and ending with 1:2 EtOAc-hexane to furnish N-[(1S)-2-butyl-1-(hydroxymethyl)hexyl]-5-chloro-2-thiophenesulfonamide as an amorphous white solid (0.630g, 16.9 %). Mass Spectrum (-ESI): 366 (M-H)⁻. Anal. Calc'd for C₁₅ H₂₅ N Cl O₃ S₂: C, 48.96; H, 7.12; N, 3.81. Found: C, 49.08, H, 6.83, N, 3.82.

Example 206

N-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide.



A. 2-Furansulfonyl chloride

5 Phosphorus pentachloride (15.29g, 73.44 mmol) was added portionwise (caution, foaming) over 5 min at 0°C to chlorosulfonic acid (21.39g, 183.6 mmol) and the resulting solution was stirred at 0°C for 10 min. Then, furan (5.0 g, 73.44 mmol) was added in one portion and the resulting dark suspension was stirred at 0°C for 15 min during which time foaming occurred and subsided. The
10 reaction mixture was then poured onto ice and the resulting suspension was extracted with CH₂Cl₂ (150 mL). The organic extract was filtered through a pad of celite, washed with brine and dried over MgSO₄. The solvent was removed in vacuo to provide 2-furansulfonyl chloride as a black oil (1.01g, 7.9 %). ¹H-NMR (DMSO-d₆, 400MHz) δ 7.4 (d, 1H); 6.38 (d, 1H); 6.35 (d, 1 H).

15 B. *N*-[(1*S*,2*S*)-1-(Hydroxymethyl)-2-methylbutyl]-2-furansulfonamide


 2-Furansulfonyl chloride (1.01g, 8.69 mmol) was added dropwise (5 min) as a solution in CH₂Cl₂ (5 mL) to a 0° C solution of L-isoleucinol (0.909 g, 7.83 mmol) in CH₂Cl₂ (20 mL) and triethylamine (2.42 mL, 17.38 mmol). The solution was allowed to warm to 25°C overnight (19 h). An aliquot was taken and TLC (1:1
20 EtOAc-hexane) indicated that reaction was complete. It was diluted with CH₂Cl₂ (100 mL) and the organic layer was washed with 1N HCl (2x50 mL) and saturated aqueous NaCl (50 mL). The organic layer was dried over MgSO₄, filtered, and concentrated to obtain a crude black oil (0.65 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluent: starting with 1:3 EtOAc-hexane and

ending with 1:2 EtOAc-hexane to furnish *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide as an amorphous white solid (0.155g, 72.12 %). Mass Spectrum (-ESI): 246 (M-H)⁻. Anal. Calc'd for C₁₀H₁₇ClNO₄S: C, 48.57; H, 6.93; N, 5.66. Found: C, 48.72; H, 6.78; N, 5.39.

5

Example 207

N-[(1*S*,2*S*)-1-(Hydroxymethyl)-2-methylbutyl]-5-iodo-2-thiophenesulfonamide



- 10 A. 5-Bromo-*N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide.
- 5-Bromothiophene-2-sulfonyl chloride (5.0 g, 19.11 mmol) was added dropwise (5 min) as a solution in CH₂Cl₂ (10 mL) to a 0° C solution of L-isoleucinol (2.108g, 18.16 mmol) in CH₂Cl₂ (15 mL) and triethylamine (3.77 mL, 27.24 mmol). The solution was allowed to warm to 25°C overnight (19 h). An aliquot was taken and
- 15 TLC (1:1 EtOAc-hexane) indicated that reaction was complete. It was diluted in methylene chloride (100 mL) and the organic layer was washed with 1N HCl (2x50 mL) and saturated aqueous NaCl (50 mL). The organic layer was dried over MgSO₄, filtered, and concentrated to obtain a crude off-yellow solid (5.2 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluent:
- 20 starting with 1:4 EtOAc-hexane and ending with 1:1 EtOAc-hexane to furnish 5-bromo-*N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide as an amorphous white solid (4.3g, 70.49 %). Mass Spectrum (-ESI): 246 (M-H)⁻.


B. *N*-[(1*S*,2*S*)-1-(Hydroxymethyl)-2-methylbutyl]-5-(tributylstannyl)-2-thiophenesulfonamide

Bis(Tributyltin) (9.28 mL, 18.52 mmol) and tetrakis(triphenyl phosphine)palladium (0) (0.7133g, 0.617 mmol) were added to a solution of 5-bromo-
 5 *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide (4.2g, 12.34 mmol) in 1,4-dioxane (42 mL). The brown solution was heated to reflux overnight (19h). An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was complete. The slurry was then filtered and the solvent removed *in vacuo* to obtain a crude yellow oil (2.1 g). The crude product was purified by column chromatography,
 10 silica gel 230-400 mesh, eluent: 1:2 EtOAc-hexane to furnish *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-5-(tributylstannyl)-2-thiophene sulfonamide as a yellow oil (0.88g, 12.9 %). Mass Spectrum (-ESI): 551 (M-H)⁺.

15 C. *N*-[(1*S*,2*S*)-1-(Hydroxymethyl)-2-methylbutyl]-5-iodo-2-thiophenesulfonamide

To a solution of *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-5-(tributylstannyl)-2-thiophene sulfonamide (0.35 g, 0.633 mmol) in methanol (4 mL) was added sequentially sodium acetate (0.104g, 1.27 mmol), sodium iodide (0.190g,
 20 1.27 mmol in H₂O) and Chloramine T trihydrate (0.36g, 1.27 mmol in methanol (0.5 mL)). The light-yellow solution turned red to orange upon addition of Chloramine T. The reacted was stirred at 25°C for 2h and then quenched by addition of 1M sodium bisulfite (10 mL). After addition of H₂O (10 mL), the aqueous layer was washed with diethyl ether (3 x 50 mL). The organic layer was dried over MgSO₄, filtered,
 25 concentrated to obtain a light yellow oil (0.210g). The crude product was purified by HPLC (sil(25 x 0.46 cm); flow rate, 1.0 mL/min; eluent, 6% MTBE in CH₂Cl₂) to furnish *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-5-iodo-2-thiophenesulfonamide as a white amorphous solid (0.125g, 51.02 %). Mass Spectrum (-ESI): 388 (M-H)⁺. Anal. Calc'd for C₁₁H₁₇NO₄S₂ · 0.07 EtOAc:
 30 C, 31.58; H, 4.21; N, 3.64. Found: C, 31.22; H, 4.22; N, 3.54.

Example 208

5-Fluoro-*N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide.

5 A. *N*-[(1*S*,2*S*)-1-(Hydroxymethyl)-2-methylbutyl]-5-(trimethylstannyl)-2-thiophenesulfonamide

Hexamethylditin (5.055g, 15.43 mmol) and tetrakis(triphenyl phosphine)palladium (0) (0.7133g, 0.617 mmol) were added to a solution of 5-bromo-*N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide (prepared as
10 in Example 199, Part A) (3.5g, 10.27 mmol) in 1,4-dioxane (70 mL). The brown solution was heated to reflux overnight (19h). An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was complete. The slurry was then filtered and the solvent removed *in vacuo* to obtain a crude yellow oil (2.1 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluent: 1:2 EtOAc-hexane to obtain *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-5-(trimethylstannyl)-
15 2-thiophenesulfonamide as a yellow oil (3.1g, 70.8%). Mass Spectrum (-ESI): 425 (M-H)⁺. Anal. Calc'd for C₁₁H₁₇NiO₄S₂: C, 36.64; H, 5.91; N, 3.29. Found: C, 36.64; H, 5.81; N, 3.21.

20 B. 5-Fluoro-*N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide


A solution of *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-5-(trimethylstannyl)-2-thiophene sulfonamide (1.0g, 2.34 mmol) in dry acetonitrile (20 mL) was stirred under nitrogen at 25°C. Selectfluor (0.850g, 2.40 mL) was added in
25 one portion and the solution stirred for 19 h at 25°C. After 3 h a white precipitate began to appear. An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was not complete. Mainly, starting material was present. The reaction was

heated to 80°C for 6 h. An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was complete. The slurry was then filtered and the solvent removed *in vacuo* to obtain a crude yellow oil (0.6g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluent: 1:2 EtOAc-hexane to obtain 5-fluoro-*N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methyl-butyl]-2-thiophenesulfonamide as an amorphous white solid (0.102g, 15.49 %). Mass Spectrum (-ESI): 280 (M-H)⁺. Anal. Calc'd for C₁₀H₁₆NFO₄S₂: C, 42.69; H, 5.73; N, 4.98. Found: C, 42.47; H, 5.74; N, 4.87.

10

Example 209

4-[1-(5-Chloro-thiophene-2-sulfonylamino)-2-hydroxy-ethyl]-piperidine-1-carboxylic acid tert-butyl ester



15

A. t-Butyl 4-((1*S*)-1-(((9H-fluoren-9-ylmethoxy)carbonyl)amino)-2-hydroxyethyl)piperidine-1-carboxylate

20

To a solution of cyanuric chloride (1.44 g, 7.80 mmol) in DME (40 mL) was added *N*-methyl morpholine (0.79 g, 7.80 mmol) at 25°C. A white precipitate formed and to this mixture 4-[carboxy-(9H-fluoren-9-ylmethoxycarbonylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester (3.75 g, 7.80 mmol) was added as a solution dissolved in DME (20 mL). After 5 h, the mixture was filtered and the liquid filtrate was cooled in an ice bath to 0°C and NaBH₄ (0.44 g, 11.63 mmol) previously dissolved in H₂O (15 mL) was added by pipet. The reaction mixture was stirred for an additional 20 min at 0°C. Diethyl ether (100 mL) was added followed by acidification using 1N HCl solution. The organic phase was then separated and washed with a 10% solution of Na₂CO₃ followed by

brine and then dried over MgSO_4 . Filtration and evaporation produced a crude glass that was flash chromatographed using ethyl acetate-hexane, 1-1 as eluent. This gave the desired product as a solid (1.03 g, 28%). MS (+ESI) 367.1 ($[\text{M}+\text{H}]^+$); 282.2; 189.1.


5 B. t-Butyl 4-[(1S)-1-amino-2-hydroxyethyl]-1-piperidinecarboxylate

To tert-butyl 4-((1S)-1-[(9H-fluoren-9-ylmethoxy)carbonyl]amino)-2-hydroxyethyl)piperidine-1-carboxylate (0.95 g, 2.03 mmol) was added 20 % piperidine in dimethylformamide (20 mL) all at once. The reaction was stirred overnight at 25°C. The dimethylformamide was evaporated off and the crude residue
10 was subjected to flash chromatography using methylene chloride-methanol-ammonium hydroxide, 95-5-0.1% as eluent. This yielded the amine product as an oil that crystallized upon standing (0.392 g, 80%). MS (+ESI) 245.2 ($[\text{M}+\text{H}]^+$); 189.2; 150.2.

15 C. 4-[1-(5-Chloro-thiophene-2-sulfonylamino)-2-hydroxy-ethyl]-piperidine-1-carboxylic acid tert-butyl ester

To a stirred mixture of tert-butyl 4-[(1S)-1-amino-2-hydroxyethyl]-1-piperidinecarboxylate (0.107 g, 0.44 mmol), triethylamine (0.046 g, 0.46 mmol) and methylene chloride (5 mL) cooled to 0°C, was added 5-chlorothiophene-2- sulfonyl
20 chloride (0.095 g, 0.44 mmol) as a solution dissolved in 2 mL methylene chloride, dropwise by pipet. After 15 min, the ice bath was removed and the reaction allowed to attain 25°C and stir overnight. The reaction was quenched by pouring it into saturated sodium bicarbonate solution (25 mL) and additional methylene chloride (15 mL). The organic phase was separated and washed sequentially with 1N HCl solution,
25 H_2O , brine and dried over MgSO_4 . The organic phase was filtered and evaporated to produce a crude oil that was flash chromatographed using ethyl acetate-hexane, 1-1 as eluent. This produced the title compound as a solid (0.109 g, 58%). MS (+APCI) 442.18 ($[\text{M}+\text{NH}_4]^+$); 386.08; 357.01; 325.07; 307.01; 285.06. Anal. Calc'd for $\text{C}_{16}\text{H}_{23}\text{ClN}_2\text{O}_5\text{S}_2$: C, 45.22; H, 5.93; N, 6.59; Found: C, 45.31; H, 5.87; N, 6.44.


30

Example 210**N-[(1S,2S)-1-(Hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide**

- 5 To a solution of 2-thiophenesulfonyl chloride (1 g, 5.48 mmol) in CH_2Cl_2 (5 mL) and (S)-isoleucinol (642 mg, 5.48 mmol) was added Hunig's base (1.05 mL, 6.02 mmol). The reaction mixture was stirred at 25°C for 24 h. The solvent was removed and the oil was dissolved in EtOAc (100 mL). The solution was washed with water (2 x 100 mL), brine (1 x 100 mL), and dried over Na_2SO_4 . The desired sulfonamide (m/z = 264.0(M+H), r_t = 0.79 min) was isolated by semi-preparative RP-HPLC using the conditions outlined for example 195.
- 10

Example 211**5-Chloro-N-[(S)-2-hydroxy-1-(4-benzylaminocyclohexyl)ethyl]thiophene-2-sulfonamide**

15



A. Part 1

A solution of (*S*)-2-(5-chiophenesulfonamido)-2-(4-hydroxycyclohexyl)-*N,O*-acetone (4.8 g, 12.7 mmol, see example 199 part 1-3) in CH₂Cl₂ (50 mL) was added to a slurry of PCC (5.46 g, 25.3 mmol), silica gel (5.46 g) and sodium acetate (1 g, 12.2 mmol) in CH₂Cl₂ (30 mL). The resulting reaction mixture was stirred at 25°C overnight. The mixture was diluted with Et₂O and filtered. The solid was washed with diethylether (3 x 50 mL) and the combined organic extracts were dried over MgSO₄. The solvent was removed *in vacuo*, and the residue was purified by column chromatography using 1:1 EtOAc/Hexane as the eluent to give the ketone as a white solid (4 g, 84%) (100% purity).

B. Part 2

To a solution of (*S*)-2-(5-chiophenesulfonamido)-2-(4-cyclohexanone)-*N,O*-acetone (340 mg, 0.9 mmol) in 1,2-dichloroethane (6 mL) was added benzylamine (118 µL, 1.08 mmol), sodium triacetoxymorohydrate (286 mg, 1.35 mmol), and acetic acid (52 µL, 0.9 mmol). The reaction was stirred at 25°C overnight whereupon the reaction was quenched with aqueous NaHCO₃ and extracted with diethyl ether and evaporated. To the resulting residue was slowly added acetic acid (10 mL of 80%) and the reaction was heated at 40 °C for nine days. The acetic acid was removed and the residue purified by column chromatography (MeOH/CH₂Cl₂/0.5-1% NH₄OH) to furnish the desired compound (254 mg, 66%) as a mixture of diastereomers.

The following compounds (Examples 211-220, Table 20) were prepared using (*S*)-2-(5-chiophenesulfonamido)-2-(4-cyclohexanone)-*N,O*-acetone (from Example 211, part 1) with benzylamine, methylamine, ethylamine, propylamine, allylamine, 3-(aminomethyl)pyridine, morpholine, 4-(aminomethyl)pyridine, 2-(aminomethyl)pyridine, and glycine ethyl ester as outlined in Example 211.




Table 20: (LCMS Data³: Molecular ion and retention time)


R-NH ₂	
benzylamine	Ex. 211 427.2 (M-H), 0.62 min 427.2 (M-H), 0.66 min
methylamine	Ex. 212 351.2 (M-H), 0.39 min 351.1 (M-H), 0.45 min
ethylamine	Example 213 365.2 (M-H), 0.47 min 365.2 (M-H), 0.53 min
n-propylamine	Ex. 214 379.2 (M-H), 0.55 min 379.2 (M-H), 0.59 min
allylamine	Ex. 215 376.9 (M-H), 0.42 min 376.9 (M-H), 0.47 min
3-(aminomethyl) pyridine	Ex. 216 427.9 (M-H), 0.35 min 427.9 (M-H), 0.41 min
morpholine	Ex. 217 406.9 (M-H), 0.40 min 406.9 (M-H), 0.44 min
4-(aminomethyl) pyridine	Ex. 218 428.0 (M-H), 0.49 min 428.0 (M-H), 0.51 min
2-(aminomethyl) pyridine	Ex. 219 428.0 (M-H), 0.36 min 428.0 (M-H), 0.37 min
glycine ethyl ester	Ex. 220 423.0 (M-H), 0.47 min 423.0 (M-H), 0.51 min

Example 221A

Method 1

5

5-Chloro-N-[(S)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide



A. 5-(1-Ethyl-propyl)-imidazolidine-2,4-dione

Sodium cyanide (12.0 g, 244.8 mmol) and 2-ethylbutyraldehyde (10.0 mL, 81.3 mmol) were added to ammonium carbonate (25.4 g, 325.3 mmol) in H₂O (300 mL). Ethanol (300 mL) was added and salts precipitated. The reaction mixture was heated to 90 °C. After 1 h, the mixture became homogeneous and was stirred at 90 °C for 18 h. After cooling to 25°C, about 500 mL of solvent was removed *in vacuo*. Concentrated HCl was added to acidify the mixture to pH 1-2 and a precipitate formed. It was filtered and the precipitate was recrystallized from EtOAc to afford 5-(1-ethyl-propyl)-imidazolidine-2,4-dione as a white solid (12.9 g, 93%).

10 Mass Spectrum (-ESI): 169 (M-H)⁺.

B. N-[(5-Chloro-2-thienyl)sulfonyl]-3-ethylnorvaline

5-(1-Ethyl-propyl)-imidazolidine-2,4-dione (12.3 g, 72.3 mmol) was dissolved in a 150 mL solution of aqueous NaOH (11.6 g, 289.2 mmol). The solution was heated by microwave in a sealed vessel for 1 h. (Microwave conditions: 15 min @ 100 % power, 150 °C, 50 psi, then 5 min 0% power, then 15 min @ 100% power, 150 °C, 50 psi, then repeat sequence.) Water and ammonium hydroxide were removed from the reaction mixture *in vacuo* and the resulting crude amino acid and NaOH mixture was used in the next reaction without further purification.

20 The crude amino acid and NaOH mixture was dissolved in 300 mL of water. The mixture was cooled to 0 °C in an ice bath. 5-Chlorothiophene-2-sulfonyl chloride (17.3 g, 79.5 mmol) was dissolved in 100 mL of THF and added dropwise to the reaction mixture over 0.5 h. After 1 h the reaction mixture was allowed to warm gradually to 25°C and stirred for 16 h. THF was removed *in vacuo* and then the mixture was acidified to pH 1 with 1N HCl. After about 15 min, a precipitate began to crash out of the milky white solution. After 1 h, the mixture was cooled in a refrigerator for 1 h and then filtered. The precipitate was washed with 1 N HCl to provide N-[(5-chloro-2-thienyl)sulfonyl]-3-ethylnorvaline as a white solid (18.5 g, 78%). Mass Spectrum (-ESI): 325 (M-H)⁺.

25

30

C. *N*-[(5-Chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline

(+)-(1*S*,2*R*)-Ephedrine hemihydrate (16.7 g, 95.6 mmol) was added to a suspension of *N*-[(5-chloro-2-thienyl)sulfonyl]-3-ethylnorvaline (31.2 g, 95.6 mmol) in 185 mL of EtOH. The mixture was heated slightly to dissolve solids and a precipitate formed. After cooling at 5 °C for 18 h the resulting suspension was filtered and the precipitate was washed with cold EtOH and EtOAc to give 27 % yield of the diastereomeric salt. The salt was recrystallized from boiling EtOAc (420 mL), then filtered off. The resulting white solid was then dissolved in 300 mL of EtOAc and 300 mL of 1*N* HCl. The layers were separated and the organic extract was washed with 1*N* HCl (2 x 200 mL), dried (Na₂SO₄), and concentrated to give *N*-[(5-chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline as a white solid (5.6 g, 18 %). Chiral HPLC [chiralpak AD (25 x 0.46 cm), 8:2 hexane (0.1% TFA) : isopropanol, L-isomer elutes at 9.6 min and D-isomer elutes at 13.1 min] indicated 96% chiral purity. $[\alpha]_D^{25} = +44.5^\circ$ (c = 1% SOLUTION, MeOH). Mass Spectrum (-ESI): 325 (M-H)⁺. Anal. Calc'd for C₁₁H₁₆ClNO₄S₂: C, 40.55; H, 4.95; N, 4.30. Found: C, 40.30; H, 4.78; N, 4.16.

D. 5-Chloro-*N*-[(*S*)-2-ethyl-1-(hydroxymethyl)butyl]-2-thiophenesulfonamide

To *N*-[(5-Chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline (5.6 g, 17.2 mmol) in THF (150 mL) at 0 °C was added a solution of 1 M borane tetrahydrofuran complex in THF (69 mL, 69 mmol) dropwise via addition funnel. After 15 min, the reaction mixture was warmed to 25°C and stirred for 18 h. It was then quenched with 90 mL of 10% AcOH in MeOH slowly. Volatiles were removed *in vacuo*. The residue was then dissolved in EtOAc (300 mL) and washed with sat. aqueous NaHCO₃ (3 x 200 mL), dried (Na₂SO₄), and concentrated to a white precipitate (5.1 g, 96% yield, 96 % chiral purity). The precipitate was recrystallized with heptane/EtOAc, 4:1, to give optically pure 5-chloro-*N*-[(*S*)-2-ethyl-1-(hydroxymethyl)butyl]-2-thiophenesulfonamide as white needles (4.4 g, 81% yield). $[\alpha]_D^{25} = +4.5^\circ$ (c = 1% SOLUTION, DMSO). Mass Spectrum (-ESI): 310 (M-H)⁺. Anal. Calc'd for C₁₁H₁₈ClNO₃S₂: C, 42.37; H, 5.82; N, 4.49. Found: C, 42.37; H, 5.79; N, 4.38.

E. 5-Chloro-*N*-[(*S*)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide

Pyridinium dichromate (2.4 g, 6.4 mmol) was added to a solution of 5-chloro-*N*-[(*S*)-2-ethyl-1-(hydroxymethyl)butyl]-2-thiophenesulfonamide (0.5 g, 1.6 mmol) in CH₂Cl₂ (20 mL). After 18 h, the reaction mixture was filtered through a
 5 plug of Celite. The filtrate was concentrated and the resulting residue was purified by silica gel column chromatography (eluant: 1:4 EtOAc-hexane) to give 5-chloro-*N*-[(*S*)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide as a white solid (303 mg, 61%).
 $[\alpha]_D^{25} = +136.76^\circ$ (c = 1% SOLUTION, CHCl₃). Mass Spectrum (-ESI): 308 (M-H)⁻.
 Anal. Calc'd for C₁₁H₁₆ClNO₃S₂: C, 42.64; H, 5.21; N, 4.52. Found: C, 42.57; H,
 10 5.24; N, 4.52.

Example 221B

Method 2

5-Chloro-*N*-[(*S*)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide

15 A. (*S*)-3-Ethyl-2-{[(*S*)-1-phenylethyl]amino} pentanenitrile

To (*S*)-(-)- α -methylbenzylamine hydrochloride salt (1.2 g, 7.6 mmol) in 80 mL of 1:1 MeOH/H₂O was added potassium cyanide (0.5 g, 7.6 mmol) and 2-methylbutyraldehyde (0.94 mL, 7.6 mmol). A precipitate formed after 30 min. After
 20 h, the suspension was filtered and washed with H₂O to give (*S*)-3-ethyl-2-{[(*S*)-1-phenylethyl]amino} pentanenitrile as a white powder (1.29 g, 74%). Mass Spectrum (+ESI): 310 (M+H)⁺. Anal. Calc'd for C₁₅H₂₂N₂: C, 78.21; H, 9.63; N, 12.16. Found: C, 77.90; H, 9.75; N, 12.32.

B. 3-Ethyl-*N*²-[(*S*)-1-phenylethyl]-L-norvalinamide

25 To 25 mL of sulfuric acid at 0 °C was added (*S*)-3-ethyl-2-{[(*S*)-1-phenylethyl]amino} pentanenitrile (2.7 g, 11.6 mmol) portionwise. The mixture was warmed to 25°C. After 2 days, the reaction mixture was poured over about 100 g of crushed ice. Concentrated NH₄OH was added to neutralize the acid. This mixture was extracted with EtOAc (3 x 100 mL), dried over Na₂SO₄, filtered and concentrated
 30 to give 3-ethyl-*N*²-[(*S*)-1-phenylethyl]-L-norvalinamide (2.6 g, 90%), which was used in the next step without purification. Mass Spectrum (+ESI): 249 (M+H)⁺. Anal.

Calc'd for $C_{15}H_{24}N_2O$: C, 72.54; H, 9.74; N, 11.28. Found: C, 72.24; H, 10.04; N, 11.01

C. 3-Ethyl-L-norvalinamide

5 A mixture of 3-ethyl- N^2 -[(*S*)-1-phenylethyl]-L-norvalinamide (2.6 g, 10.5 mmol) and 5% Pd/C (800 mg) was shaken for 24 h in a Parr apparatus under 3 atm of H_2 . The mixture was filtered through a plug of Celite and the solvent was removed *in vacuo* to give 3-ethyl-L-norvalinamide as a white solid (1.4 g, 93 %), which was used in the next reaction without further purification. Mass Spectrum
10 (+ESI): 145 (M+H)⁺.

D. *N*-[(5-Chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline

3-Ethyl-L-norvalinamide (1.2 g, 4.8 mmol) was dissolved in conc. HCl (10 mL) and heated to 100 °C for 16 h. The reaction mixture was concentrated to a
15 white solid consisting of the amino acid hydrochloride salt and one equivalent of NH_4Cl , which was used in the next reaction without purification.

Amino acid hydrochloride salt with 1 equivalent of NH_4Cl (0.28 g, 1.19 mmol) was dissolved in 6 mL of H_2O and then NaOH (0.24 g, 6.00 mmol) was added. The solution was cooled to 0 °C and then 5-chlorothiophene-2-sulfonyl
20 chloride (0.29 g, 1.32 mmol) in 6 mL of THF was added dropwise. The mixture was warmed to 25°C. After 19 h, THF was removed *in vacuo*. The remaining solution was diluted with 10 mL of H_2O and washed with EtOAc (2 x 10 mL). The solution was acidified with 1N HCl and a precipitate formed. This was filtered to give *N*-[(5-chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline as a white solid (0.17 g, 44%). Chiral
25 HPLC indicates that only the *S* enantiomer is present.

5-Chloro-*N*-[(*S*)-2-ethyl-1-(hydroxymethyl)butyl]-2-thiophenesulfonamide and 5-Chloro-*N*-[(*S*)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide were then prepared from *N*-[(5-Chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline according to
30 method 1 of Example 221A.


Example 221C**5-Chloro-N-[(S)-2-ethyl-1-(hydroxymethyl)butyl]-2-thiophenesulfonamide**

Into a 3L 3-necked flask equipped with a nitrogen inlet tube, a mechanical stirrer, and an addition funnel with a stopper was placed lithium borohydride (145 mL of a 2 M solution in THF, 0.29 mol). The solution was placed under nitrogen and cooled to 0°C. Chlorotrimethylsilane (73.8 mL, 0.58 mol) was added dropwise over a period of 30 min. The ice bath was removed and the resulting slurry was stirred at room temperature for 30 min. The reaction mixture was cooled to 0°C and the 2-(S)-amino-3-ethyl-pentanoic acid (21.1 g, 0.145 mol), which was prepared according to Scheme 13, was added in portions as a solid over a period of 15 min. The reaction mixture was allowed to warm slowly to room temperature as the ice bath melted. After 3 days at room temperature, the reaction mixture was cooled to 0°C, and methanol (217 mL) was carefully added over a period of 80 min. The solution was stirred at room temperature for an additional 40 min, then concentrated under reduced pressure in a water bath at 60°C. The resulting slurry was made basic with 20% sodium hydroxide (37.5 mL). Water (37.5 mL) was added, and the entire aqueous layer was extracted with methylene chloride (300 mL), and dried (Na₂SO₄). Concentration under reduced pressure gave 2(S)-amino-3-ethylpentanol as an oil (17.3 g, 91%), which was used immediately or stored in the freezer overnight: Opt. Rot. [α]_D²⁵ = -3.7° (1% solution, DMSO); ¹H NMR (DMSO-d₆, 500 MHz): δ 4.38 (broad s, 1H), 3.35 (dd overlapping with a broad s at δ 3.32, J=4.5, 10.3 Hz, 3H), 3.14 (dd, J=7.9, 10.2 Hz, 1H), 2.63 (m, 1H), 1.45-1.05 (m, 5H), 0.82 and 0.81 (two overlapping triplets, J=7.4 Hz, 6H); MS(+ESI): [M+H]⁺, 132 (60%).

A mixture of 2(S)-amino-3-ethylpentanol (34.1 g, 0.26 mol) and methylene chloride (700 mL) was placed under Argon, and cooled to 0°C. Triethylamine (36.2 mL, 0.26 mol) was added, followed by the dropwise addition of 5-chlorothiophene-2-sulfonyl chloride (56.4 g, 0.26 mol) in methylene chloride (400 mL). The reaction mixture was allowed to warm slowly to room temperature as the ice bath melted. After 3 days at room temperature, the reaction mixture was divided into two 0.6 L portions. Each portion was diluted with ethyl acetate (1L), and washed three times with saturated potassium phosphate monohydrate (200 mL), once with brine (200 mL), and dried (Na₂SO₄). Concentration under reduced pressure gave a white solid

(74.5 g, 92%). The product (87.98 g) from several runs were combined and recrystallized from hot heptane:ethyl acetate (4:1, 775 mL) to give the title compound as crystals (74.9 g, 85%): mp 115-117.6°C; Opt. Rot. $[\alpha]_D^{25} = +10.81^\circ$ (1% solution, MeOH); ^1H NMR (DMSO- d_6 , 500 MHz): δ 7.71 (d, $J=8.1$ Hz, 1H), 7.44 (d, $J=4.1$ Hz, 1H), 7.22 (d, $J=4.1$ Hz, 1H), 4.56 (t, $J=5.2$ Hz, OH), 3.31-3.15 (m, 3H), 1.40-1.15 (m, 4H), 1.07 (m, 1H), 0.79 and 0.76 (two overlapping triplets, $J=7.3$ Hz, 6H); ^{13}C NMR (DMSO- d_6 , 100 MHz): δ 141.75, 133.73, 130.95, 127.60, 60.41, 56.89, 41.57, 21.31, 20.80, 11.79, 11.51; MS(-ESI): $[\text{M}-\text{H}]^-$, 1 chlorine isotope pattern, 310 (100%), 312 (30%); Anal. Calc. for $\text{C}_{11}\text{H}_{18}\text{ClNO}_3\text{S}_2$: C, 42.37, H, 5.82, N, 4.49. Found: C, 42.34, H, 5.65, N, 4.43. Chiral HPLC (Chiralpak AD, 25x0.46 cm, eluant 8:2 hexane/isopropanol containing 0.1% TFA, flow rate 0.5 mL/min, UV detection at 254 nm, retention times for the S and R isomers are 10.95 min and 11.95 min, respectively) revealed an S/R ratio of 100.0:0.0.

15


Example 222**5-Chloro-*N*-[(*S*)-2-ethyl-1-(1-hydroxyethyl)butyl]thiophene-2-sulfonamide**

A solution of methylmagnesium bromide (1.4 M, 7.0 mL, 9.7 mmol) in toluene/THF (75:25) was added to a 0 °C solution of 5-chloro-*N*-[(*S*)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide (Example 221, 1.0 g, 3.2 mmol) in THF (30 mL). The mixture was warmed to 25°C and after 2 h was quenched carefully with saturated aqueous ammonium chloride (25 mL). The mixture was extracted with EtOAc (3 x 25 mL). The organic extract was dried over Na_2SO_4 , filtered and concentrated to give a colorless oil. The product was purified by column chromatography (Biotage), eluant: 1:4 EtOAc-hexane, to afford 5-chloro-*N*-[(*S*)-2-ethyl-1-(1-hydroxyethyl)butyl]thiophene-2-sulfonamide as a white solid (876 mg, 83%). The product is a diastereomeric mixture with a ratio of 3:7. mp 95-98 °C.

Anal. Calc'd for $C_{12}H_{20}ClNO_3S_2$: C, 44.23; H, 6.19; N, 4.30. Found: C, 44.25; H, 6.35; N, 4.29. Mass Spectrum (-ESI): 324 (M-H)⁺.

Example 223

5 5-Chloro-*N*-[(*S*)-2-ethyl-1-(1-hydroxy-1-methylethyl)butyl]thiophene-2-sulfonamide



10 A. *N*-[(5-Chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline methyl ester
Trimethylsilyldiazomethane (3.1 mL, 6.1 mmol) was added to a solution of *N*-[(5-chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline (1.0 g, 3.1 mmol) in THF (20 mL) and MeOH (5 mL). After 2 h the mixture was concentrated to give *N*-[(5-chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline methyl ester as a white solid (1.0 g,
15 99%). Mass Spectrum (-ESI): 338.00 (M-H)⁺.


B. 5-Chloro-*N*-[(*S*)-2-ethyl-1-(1-hydroxy-1-methylethyl)butyl]thiophene-2-sulfonamide

A solution of methylmagnesium bromide (1.4 M, 9.5 mL, 13.2 mmol)
20 in toluene/THF (75:25) was added to a 0 °C solution of *N*-[(5-chloro-2-thienyl)sulfonyl]-3-ethyl-L-norvaline methyl ester (0.90 g, 2.65 mmol) in THF (26 mL). The solution was allowed to warm to 25 °C, then heated to 55 °C and stirred for 18 h. It was then cooled to 0 °C and quenched slowly with saturated aqueous NH_4Cl . EtOAc (75 mL) was added and the phases were separated. The organic layer was
25 dried over Na_2SO_4 , filtered and concentrated to give a yellow oil. The crude product was purified by column chromatography (Biotage), eluant: 1:4 EtOAc-hexane, to afford 5-chloro-*N*-[(*S*)-2-ethyl-1-(1-hydroxy-1-methylethyl)butyl]thiophene-2-sulfonamide as a colorless oil (0.72 g, 80%). Mass Spectrum (-ESI): 338 (M-H)⁺.

Anal. Calc'd for $C_{13}H_{22}ClNO_3S_2$: C, 45.94; H, 6.52; N, 4.12. Found: C, 46.10; H, 6.63; N, 4.04.

Example 224

5 **5-Chloro-*N*-(2-hydroxy-1-tetrahydro-*H*-thiopyran-4-ylethyl)thiophene-2-sulfonamide**



10 A. (5-Chloro-thiophene-2-sulfonylamino)-(tetrahydro-thiopyran-4-yl)-acetic acid

Sodium hydroxide (0.20 g, 5.04 mmol) was added to a mixture of *N*-Fmoc-amino-(4-tetrahydrothiopyranyl)acetic acid (0.50 g, 1.26 mmol) in MeOH:water, 2:1 (15 mL) at 25 °C. The reaction mixture was allowed to stir for 20 h. TLC (1:9 MeOH/CHCl₃) indicated that the reaction was complete. The mixture was diluted with water and washed with EtOAc. The water layer was concentrated to give a white solid with NaOH remaining. This white solid was redissolved in H₂O:THF, 1:2 (15 mL) and cooled to 0 °C. 5-chlorothiophene-2-sulfonyl chloride (0.683 g, 3.15 mmol) was dissolved in THF (2 mL) and added to the mixture dropwise which was then warmed to 25°C overnight. Aqueous 1 N HCl was added to acidify the mixture to pH 1. EtOAc was added and the layers were separated. The organic extract was washed with 1 N HCl, and H₂O, dried over Na₂SO₄, filtered and concentrated to give (5-chloro-thiophene-2-sulfonylamino)-(tetrahydro-thiopyran-4-yl)-acetic acid as a reddish black solid (0.14, 31%), which was used in the next reaction without purification. Mass Spectrum (+ESI): 357 (M+H)⁺


25

B. 5-Chloro-*N*-(2-hydroxy-1-tetrahydro-*H*-thiopyran-4-ylethyl)thiophene-2-sulfonamide

(5-Chloro-thiophene-2-sulfonylamino)-(tetrahydro-thiopyran-4-yl)-
5 acetic acid (0.14 g, 0.40 mmol) was dissolved in THF (2 mL) and cooled to 0 °C. A
solution of borane tetrahydrofuran complex (1 M, 3.2 mL, 3.2 mmol) in THF was
added dropwise and the mixture was allowed to warm to 25 °C overnight. The volatile
solvents were removed in vacuo and the resulting orange oil was diluted with EtOAc
and washed with H₂O, 1 N HCl, and saturated aqueous NaHCO₃. The organic extract
10 was dried over Na₂SO₄, filtered and concentrated. The resulting residue was purified
by column chromatography (Biotage), eluant: 1:1 EtOAc:hexane, to afford 5-chloro-
N-(2-hydroxy-1-tetrahydro-*H*-thiopyran-4-ylethyl)thiophene-2-sulfonamide (40 mg,
30%) as a white solid. mp 108-110 °C. Mass Spectrum (-ESI): 340 (M-H)⁻. Anal.
Calc'd for C₁₁H₁₆ClNO₃S₃: C, 38.64; H, 4.72; N, 4.10. Found: C, 38.80; H, 4.69; N,
15 3.88.


Example 225

5-Chloro-*N*-[(*S*)-2-hydroxy-1-piperidin-4-ylethyl]thiophene-2-sulfonamide



20 To a solution of 4-[1-(5-chloro-thiophene-2-sulfonylamino)-2-hydroxy-ethyl]-
piperidine-1-carboxylic acid tert-butyl ester (0.204 g, 0.48 mmol (see example 209))
dissolved in dichloromethane (2 mL) at 0°C was added trifluoroacetic acid (0.5 mL).
The reaction was allowed to warm to 25°C and stir overnight. The mixture was then
concentrated and dichloromethane was added and evaporated 6 times to yield a crude
25 solid. Purification by HPLC (C-18 column, (21x75 mm) with elution system 60-100%
acetonitrile-water+0.1% TFA, 20 min gradient) gave the product as an oil (0.0166 g,
11%). MS (ESI) *m/z* 325 ([M+H]⁺).

Example 226

N-[(*S*)-2-Ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide


- 5 A. *N*-[(1*S*)-2-Ethyl-1-(hydroxymethyl)butyl]-5-(trimethylstannyl)thiophene-2-sulfonamide.

A solution of 5-bromo-*N*-[(1*S*)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide (0.71 g, 2.0 mmol), hexamethylditin (0.983 g, 3.0 mmol),
 10 tetrakis(triphenylphosphine)palladium (0.115 g, 0.10 mmol) and 1,4-dioxane (15 mL) was refluxed for 16 h under a nitrogen atmosphere. After cooling to 25°C, dichloromethane (10 mL) was added and the mixture filtered and evaporated to give the product as a crude oil (0.49g), that was used without purification in the next step, part B. MS (-ESI) 439.20 ([*M*-H]⁻).

15

- B. *N*-[(*S*)-2-Ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide


To a stirred mixture of anhydrous acetonitrile (6 mL) and *N*-[(1*S*)-2-ethyl-1-(hydroxymethyl)butyl]-5-(trimethylstannyl)thiophene-2-sulfonamide (0.24 g, 0.56 mmol) was added Selectfluor (Aldrich) (0.204 g, 0.57 mmol) all at once. The
 20 mixture was heated to 75°C under a nitrogen atmosphere, stirred 16 h and then cooled to 25°C and filtered. Evaporation of the solvent produced a crude solid that was taken up in ethyl acetate and again filtered to remove insoluble solids. Evaporation of the remaining solvent produced an oil that was purified by flash chromatography using hexane-ethyl acetate 2-1 as eluant, producing the title compound as the major product
 25 (0.051 g, 33%). MS (-ESI) 276.20 ([*M*-H]⁻).

Example 227***N*-[(*S*)-2-Ethyl-1-(hydroxymethyl)butyl]-5-fluorothiophene-2-sulfonamide**

- 5 This compound was synthesized as a side product using the procedure found in example 226 (parts A and B) and was isolated from the same flash chromatography column as a solid (0.024 g, 15%). MS (-ESI) 294.20 ([M-H]⁻).

Example 228

10 **5-Chloro-N-[(1*S*)-1-(2,3-dihydro-1*H*-inden-2-yl)-2-hydroxyethyl]thiophene-2-sulfonamide**



A. 9*H*-Fluoren-9-ylmethyl-(1*S*)-1-(2,3-dihydro-1*H*-inden-2-yl)-2-hydroxyethylcarbamate

- 15 1*N* Borane-THF (24.18 mL) was added dropwise over 30 min at 0° C to a solution of (2*S*-2,3-dihydro-1*H*-indene-2-yl)[[(9*H*-fluoren-9-ylmethoxy)carbonyl]amino]ethanoic acid (2.0 g, 4.84 mmol) in anhydrous tetrahydrofuran (20 mL). The reaction was allowed to warm to 25°C overnight, and then was quenched by addition of 10.0 mL of 10% acetic acid in methanol. After
- 20 solvent evaporation, the crude product was dissolved in ethyl acetate and washed with 1*N* HCl, water and 10% NaHCO₃. The organic layer was dried over MgSO₄, filtered and concentrated to obtain a crude yellow oil (1.8 g). The crude product was purified


by column chromatography, silica gel 230-400 mesh, eluant: 1:2 EtOAc-hexane. This produced the title compound as an amorphous solid (1.05 g, 54.4 %). Mass Spectrum. (-ESI): 398 (M-H)⁻. (+ESI): 400 (M+H)⁺.

5 B. (2S)-2-Amino-2-(2,3-dihydro-1H-inden-2-yl)ethanol


20 % Piperidine in DMF (15 mL) was added to a solution of 9H-fluoren-9-ylmethyl-(1S)-1-(2,3-dihydro-1H-inden-2-yl)-2-hydroxyethylcarbamate (1.05 g, 2.63 mmol) in DMF (5 mL). The reaction was stirred at 25°C for 19 h. After solvent evaporation, the crude product was dissolved in ethyl acetate (50 mL) and
10 dried over MgSO₄, filtered and concentrated to obtain a crude yellow oil (1.05 g). Mass Spectrum (+ESI): 179 (M+H)⁺.

 C. 5-Chloro-N-[(1S)-1-(2,3-dihydro-1H-inden-2-yl)-2-hydroxyethyl]thiophene-2-sulfonamide

15 5-Chlorothiophene-2-sulfonyl chloride (0.856 g, 3.94 mmol) was added dropwise (5 min) as a solution in CH₂Cl₂ (5 mL) to a 0°C solution of (2S)-2-amino-2-(2,3-dihydro-1H-inden-2-yl)ethanol (0.46 g, 2.63 mmol) in CH₂Cl₂ (5 mL) and triethylamine (3.8 mL, 5.26 mmol). The solution was allowed to warm to 25°C overnight (19 h). An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that
20 reaction was complete. It was diluted with CH₂Cl₂ (50 mL) and the organic layer was washed with 1N HCl (2x50 mL), saturated aqueous NaCl (50 mL). The organic layer was dried over MgSO₄, filtered and concentrated to obtain a crude oil (0.89 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluant: 1:4 EtOAc-hexane to afford 5-chloro-N-[(1S)-1-(2,3-dihydro-1H-inden-2-yl)-
25 2-hydroxy ethyl]thiophene-2-sulfonamide as an amorphous white solid (0.361 g, 38.4 %). Mass Spectrum (-ESI): 356 (M-H)⁻. Anal. Calc'd for C₁₅H₁₀ClNO₃S₂: C: 50.34 H: 4.51 N: 3.91 Found: C: 50.28 H: 4.36 N: 3.77.

Example 229**5-Chloro-N-[(1S,2S)-1-[(Z)-(hydroxyimino)methyl]-2-methylbutyl]thiophene-2-sulfonamide**

- 5 A solution of 5-chloro-N-[(1S,2S)-1-formyl-2-methylbutyl]thiophene-2-sulfonamide (Example 118, 1.0 g, 3.4 mmol), hydroxylamine hydrochloride (0.464 g, 6.78 mmol) and sodium acetate (0.556 g, 6.78 mmol) in methanol (10 mL) was stirred under reflux for 19 h. After evaporation of the solvent, the residue was diluted with aqueous K₂CO₃ (20 mL) and then extracted with CH₂Cl₂ (2x40 mL). The combined
- 10 reaction extracts were washed with brine, dried over MgSO₄, filtered and concentrated to obtain a crude oil (0.89 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluant: 1:4 EtOAc-hexane. This produced the title compound (Z-isomer) as an amorphous white solid (32 mg, 3.1 %).
- Mass Spectrum (-ESI): 309 (M-H)⁺. Anal. Calc'd for C₁₀H₁₅ClN₂O₃S₂ · 0.10 C₄H₈O₂:
 15 C: 39.08 H: 4.98 N: 8.76 Found: C: 38.72 H: 4.67 N: 8.43.

Example 230**5-chloro-N-[(S,S)-1-[(E)-(hydroxyimino)methyl]-2-methylbutyl]thiophene-2-sulfonamide**

A solution of 5-chloro-N-[(1R,2S)-1-formyl-2-methylbutyl]thiophene-2-sulfonamide (Example 118, 1.0 g, 3.4 mmol), hydroxylamine hydrochloride (0.464 g, 6.78 mmol) and sodium acetate (0.556 g, 6.78 mmol) in methanol (10 mL) was stirred under reflux for 19 h. After evaporation of the solvent, the residue was diluted with aqueous K₂CO₃ (20 mL) and then extracted with CH₂Cl₂ (2x40 mL). The combined reaction extracts were washed with brine, dried over MgSO₄, filtered, and concentrated to obtain a crude oil (0.89 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluant: 1:4 EtOAc-hexane This produced the title compound (E-isomer) as an amorphous white solid (300 mg, 28.3 %).

Mass Spectrum (-ESI): 309 (M-H)⁻. Anal. Calc'd for C₁₀H₁₅ClN₂O₃S₂ · 0.40 C₄H₈O₂: C : 40.26 H: 5.30 N: 8.09 Found: C: 39.78 H: 5.23 N: 7.77

A. Diethyl-3-ethyl-5-oxopyrrolidine-2,2-dicarboxylate

To 150 mg of sodium dissolved in 150 mL of absolute ethanol was added diethyl acetamidomalonate (5.3 g, 25 mmol) and ethyl-(2E)-pent-2-enoate (3.5 g, 27.3 mmol). The reaction mixture was then refluxed for 20 h. After this period of time, 2 mL of glacial acetic acid was added, volatiles were removed under pressure with the aid of a water aspirator and heating bath. On cooling the residue solidified. The residue was dissolved in 50 mL of toluene and to this was added 20 mL of petroleum ether. The product precipitated when the mixture was cooled. The crystals were collected and washed with water and further dried *in vacuo* to obtain a white solid (5.6 g, 79.77 %). Mass Spectrum (+ESI): 258 (M+H)⁺.

B. 3-Ethylglutamic acid.

5.6 g of Diethyl-3-ethyl-5-oxopyrrolidine-2,2-dicarboxylate (21.76, 84.67 mmol) was refluxed in 80 mL of 49 % fuming HBr for 4 h. After this time, the contents were placed in vacuo and the volatile constituents were removed. The gummy residue was dissolved in 25 mL of distilled water and the water was removed as before. The process was repeated once more. The residue was dissolved in 20 mL of water and the pH of the solution was adjusted to pH 3 with concentrated ammonia (2 mL) solution. At this point precipitation of the ethyl glutamic acid was encouraged by cooling on an ice bath or by diluting the aqueous solution with 100 mL of absolute

ethanol. Precipitation from the water-ethanol mixture is complete in 48 h. Care must be taken to add the ethanol slowly to prevent the precipitation of an undesired side product. The compound was purified by crystallization from water-ethanol (1:1) mixture. This produced the title compound as an amorphous white solid (3.5 g, 99 %).

5 Mass Spectrum (+ESI): 176 (M+H)⁺.

C. 3-Ethyl-2-methylpentane-1,5-diol

To a slurry of LAH (2.06 g, 54.29 mmol) in THF (60 mL) was added 3-ethylglutamic acid (3.5 g, 21.71 mmol) dropwise at 0°C over 20 min. The reaction
10 was heated to 36°C for 18 h. The reaction slurry (gray) was cooled to 0°C and quenched with H₂O (3 mL) then washed with 1N NaOH (9 mL) and H₂O (3 mL). It was then stirred for 6 h at 25°C to obtain an off-white slurry. The slurry was filtered and the mother liquor was further dried over MgSO₄, filtered and concentrated *in vacuo* to obtain 3-ethyl-2-methylpentane-1,5-diol as a crude yellow oil (2.85 g,
15 89.17 %). Mass Spectrum (+ESI): 170 (M+Na)⁺.

D. 5-Chloro-N-[2-ethyl-4-hydroxy-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide

5-Chlorothiophene-2-sulfonyl chloride (6.4 g, 24.48 mmol) was added dropwise (5 min) as a solution in CH₂Cl₂ (5 mL) to a 0°C solution of 3-ethyl-2-methylpentane-1,5-diol (2.85 g, 19.34 mmol) in CH₂Cl₂ (30 mL) and triethylamine (5.66 mL, 40.81 mmol). The solution was allowed to warm to 25°C overnight (19 h). An aliquot was taken and TLC (1:1 EtOAc-hexane) indicated that reaction was complete. It was diluted with CH₂Cl₂ (50 mL) and the organic layer was washed with
25 1N HCl (2x50 mL) and saturated aqueous NaCl (50 mL). The organic layer was dried over MgSO₄, filtered and concentrated to obtain a crude oil (4.9 g). The crude product was purified by column chromatography, silica gel 230-400 mesh, eluant: 1:4 EtOAc-hexane to afford 5-chloro-N-[2-ethyl-4-hydroxy-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide as an amorphous white solid (0.450 g, 7.3 %). Mass
30 Spectrum (-ESI): 326 (M-H)⁻.

1. Semi-preparative RP-HPLC conditions:

Gilson Semi-Preparative HPLC system with Unipoint Software.

Column: Phenomenex C18 Luna 21.6 mm x 60 mm, 5 μ

Solvent A: Water (0.02% TFA buffer)

5 Solvent B: Acetonitrile (0.02 % TFA buffer)

Solvent Gradient: Time 0 : 10 % B; 2.5 min : 10 % B; 14 min : 90 % B.

Flow Rate: 22.5 mL/min

The product peak was collected based on UV absorption and concentrated.

10 **2. Analytical LCMS conditions:**

Hewlett Packard 1100 MSD with ChemStation Software

Column: YMC ODS-AM 2.0 mm x 50 mm 5 μ column at 23°C;

3 μ L injection;

Solvent A: Water (0.02% TFA buffer)

15 Solvent B: Acetonitrile (0.02 % TFA buffer)

Gradient: Time 0: 95% A; 0.3 min: 95% A; 4.7 min: 10% A; 4.9 min: 95% A.

Flow rate 1.5 mL/min;

Detection: 254 nm DAD;

API-ES Scanning Mode Positive 150-700; Fragmentor 70 mV.

20

3. Analytical LCMS conditions:

ZMD (Waters) or Platform (Micromass) or LCZ (Micromass)

Column: Zorbax SB-C8

Solvent: Acetonitrile+H₂O containing 0.1%TFA or 0.1%FA

25 Gradient: Gradient: 2.5min 15% Acetonitrile – 95% Acetonitrile

Flow rate 3mL/min

Detection: ELSD detection (SEDEX 55)

UV 253 detection (Schimadzu)

30

Example 233 - Repressor Release Assay (RRA)

The compounds generated as described in Examples 1 through 220 were tested in the RRA in accordance with published techniques [Shuey, D.J., Sheiffle, P., Jones, D., Cockett, M.I., and Quinet, E.M. (1999), "Repressor release: a useful tool for
5 monitoring amyloid precursor protein (APP) proteolysis in mammalian cells", *Society for Neuroscience Abstracts*, Vol. 25, 29th Annual Meeting of Society for Neuroscience, Miami Beach, Florida, October 23-28, 1999]. Briefly, this assay is performed as follows.

A. Cell Culture

10 CHO-K1 cells are cultured in whole DMEM media (DMEM - High Glucose with 10% fetal bovine serum, 1% Non-essential Amino Acids, and 1% Penicillin-Streptomycin) at 37^o C with 5% CO₂. Two million cells are plated into 10-cm dishes 24 hrs prior to transfection.

Transient transfections are completed as recommended by Gibco BRL
15 using their Lipofectamine Plus system. First, 6 µg of pRSVO-luc and 6 µg of APP-lacI construct DNA are added to 460 µL Opti-Mem transfection media and incubated with 30 µL Plus reagent for 15 minutes. Then, a lipid mixture of 40 µL Lipofectamine reagent and 460 µL Opti-Mem transfection media is incubated with the DNA-Plus reagent mixture for 15 minutes. During the DNA-lipid incubation, the
20 CHO-K1 cells are washed once and covered in 5.0 mL DMEM media without Penicillin-Streptomycin. The DNA-lipid preparation is then layered onto these cells and incubated at 37°C overnight.

One and one half million transfected cells per well (100 µL total volume) are plated into sterile, opaque Packard 96-well Cultur-Plates in clear DMEM
25 whole media (DMEM - without phenol red) and incubated at 37^o C with 5% CO₂ for 3-5 hours.

B. Compound Dilution

Compounds are diluted using two different protocols; one protocol is used for compounds supplied neat (weighed powder in vial) and the other protocol is
30 used for compounds supplied in solution (20 mM in DMSO in 96-well plates). For both protocols, 25 mM Hepes and 25 mM Hepes/1% DMSO are prepared fresh to be

used as diluent. The Hepes/DMSO is used as the diluent control on all experimental plates.

The following table depicts the steps for compound dilution (please note that the last step is the addition of compound to cells/media in tissue culture plate):

Table 21

	Concentration	Dilution
Stock Solution	10 mg/mL	x mg compound (vial) diluted with 100% DMSO
Dilution 1	1 mg/mL	20 μ L stock solution 180 μ L 25mM Hepes
Dilution 2	200 μ g/mL	60 μ L Dilution 1 240 μ L 25 mM Hepes
Dilution 3 (in Cell Plate)	20 μ g/mL	11.3 μ L Dilution 2 (in 100 μ L cells/well)

Because some compounds arrive in 96-well format at 20 mM, the following represents the protocol for their dilution (note that an average molecular weight of these compounds was used to calculate these dilutions and as above, the last step is the addition of compound to cells/media in tissue culture plate):

Table 22

	Concentration	Dilution
Stock Solution (original conc.)	-	20 mM Solution
Dilution 1	~200 μ g/mL	6 μ L stock solution 194 μ L 25 mM Hepes
Dilution 2 (in Cell Plate)	~20 μ g/mL	11.3 μ L Dilution 2 (in 100 μ L cells/well)

Once compounds are diluted, they are applied in duplicate on cells in tissue culture plates (prepared above). Cells are incubated with compound at 37° C with 5% CO₂ for an additional 36-48 hours.

C. Assay Measurement

Luciferase assays (LucLite reagent, Packard) are performed and are read on a Packard TopCount instrument. Media is removed from each 96-well plate

and replaced with 100 μ L PBS per well (with Mg^{2+} and Ca^{2+}). An equal volume (100 μ L) of the LucLite lysis/substrate buffer is added to each well and the plates are sealed and mixed in the dark on a rotary shaker for 15-30 minutes at room temperature. Luciferase readings are then taken on the TopCount instrument.

- 5 Measurements are expressed as relative light units (RLU) and are calculated and analyzed in MS Excel as follows.

D. Analysis of data

- The results of the assay with respect to the compounds exemplified herein are provided in the following table. A compound is considered active in RRA if it leads to at least a 1.5 fold increase in luciferase activity at 20 μ M and is non-toxic, as determined by loss of signal (≤ 0.75 fold increase). Fold increase is the amount of luciferase activity (measured in relative light units) over diluent control. SEM represents the standard error of the mean for fold increase (not shown). All compounds tested were found to be non-toxic.

Table 23

Ex #	Conc (μ g/mL)	APPI Fold Increase	Name
1	10	1.57	3-bromo-5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
2	10	3.2	5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
	3	2.34	
	10	3.06	
	10	4.26	
	1	1.79	
	20	4.5	
	20	5.5	
	20	6.3	
	3	2.96	
	1	1.54	
3	10	1	4-bromo-5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
	20	1.5	
	20	25.4	

4	10	3.3	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
	3	2.98	
	3	3.52	
	20	4	
	20	4	
	20	5.4	
	20	3.3	
	20	5.1	
	1	2.22	
	10	5.03	
5	3	1.1	2,5-dichloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-3-sulfonamide
	1	0.9	
	0.3	0.9	
	10	1.1	
6	10	1.1	4,5-dichloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
	20	1.5	
	20	2.9	
7	20	3.2	N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
8	10	3.7	5-chloro-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide
	3	2.5	
	20	3.3	
	20	5	
	20	3.8	
	1	1.5	
9	10	1.81	5-bromo-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide
	20	4.1	
10	20	1.2	5-bromo-N-[(1R)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide
	20	6.4	
	20	1.7	
11	20	2	4,5-dibromo-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide
	20	4.9	
12	3	3.07	5-chloro-N-[(1S)-1-cyclohexyl-2-hydroxyethyl]thiophene-2-sulfonamide
	1	2.58	
	10	4.2	
	20	4.3	
	3	3.1	
	20	8.6	
	1	2	
	10	3.9	
13	10	3.94	5-bromo-N-[(1S)-1-cyclohexyl-2-hydroxyethyl]thiophene-2-sulfonamide
	1	2.17	
	3	4.02	
14	10	3.45	5-chloro-N-[1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide
	3	1.87	
	3	3.33	

15	10	3.08	5-bromo-N-[1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide
16	10	4.19	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
	0.3	1.7	
	1	1.9	
	3	3.1	
	20	3.7	
	20	14.2	
	20	4.8	
	20	3.2	
	20	5.6	
	20	6.1	
	10	3.2	
17	1	2.38	5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
	3	3.47	
	10	3.29	
	20	6.1	
	20	3.5	
18	10	5.23	5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide 1,1-dioxide
	20	5.7	
	3	2.19	
	20	3	
19	20	6	5-chloro-N-[1-(hydroxymethyl)-2,3-dimethylpentyl]thiophene-2-sulfonamide
20	20	5	5-chloro-N-[1-(hydroxymethyl)-2-methylpentyl]thiophene-2-sulfonamide
21	20	3.3	5-chloro-N-[2-ethyl-1-(hydroxymethyl)hexyl]thiophene-2-sulfonamide
22	20	7	5-chloro-N-[2-hydroxy-1-(2,4,6-trimethylcyclohex-3-en-1-yl)ethyl]thiophene-2-sulfonamide
23	20	6.4	5-chloro-N-(1-cyclohex-3-en-1-yl-2-hydroxyethyl)thiophene-2-sulfonamide
24	20	6	5-chloro-N-(1-cyclopentyl-2-hydroxyethyl)thiophene-2-sulfonamide
25	20	7.5	5-bromo-N-[(1S)-1-(hydroxymethyl)-1,2-dimethylpropyl]thiophene-2-sulfonamide
26	20	5	5-chloro-N-[(1S)-1-(hydroxymethyl)-1,2-dimethylpropyl]thiophene-2-sulfonamide
27	20	8.7	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2,4-dimethylpentyl]thiophene-2-sulfonamide
28	20	7	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)propyl]thiophene-2-sulfonamide
29	20	2.6	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-methyloctyl]thiophene-2-sulfonamide

30	20	6.7	5-chloro-N-[(1S)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide
	20	4.4	
31	20	4.4	5-chloro-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide
32	20	5	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)butyl]thiophene-2-sulfonamide
33	20	6.5	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylpentyl]thiophene-2-sulfonamide
34	20	5.9	5-chloro-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide
35	20	3.7	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-4-methyl-2-propylpentyl]thiophene-2-sulfonamide
36	20	3.8	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)pentyl]thiophene-2-sulfonamide
37	20	4.8	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-propyloctyl]thiophene-2-sulfonamide
38	20	4.1	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-phenylpentyl]thiophene-2-sulfonamide
39	20	5.7	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylheptyl]thiophene-2-sulfonamide
40	20	3.9	5-chloro-N-[(1S)-2-propyl-1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide
41	20	4.7	5-chloro-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)heptyl]thiophene-2-sulfonamide
42	20	3.4	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-isobutylheptyl]thiophene-2-sulfonamide
43	20	1.7	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)heptyl]thiophene-2-sulfonamide
44	20	1.6	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-pentyloctyl]thiophene-2-sulfonamide
45	20	2.1	5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-phenylheptyl]thiophene-2-sulfonamide
46	20	5.8	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide
47	20	5.4	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-4-methyl-2-phenylpentyl]thiophene-2-sulfonamide
48	20	4.7	5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide
49	20	6.1	5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide
50	20	7.7	5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide
51	20	4	5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide

52	20	7	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-isopropyloctyl]thiophene-2-sulfonamide
53	20	3.9	N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)propyl]-5-chlorothiophene-2-sulfonamide
54	20	3.3	N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)butyl]-5-chlorothiophene-2-sulfonamide
55	20	1.9	N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)-4-methylpentyl]-5-chlorothiophene-2-sulfonamide
56	20	6.8	5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2,4-dimethylpentyl]thiophene-2-sulfonamide
57	20	6.4	5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-methyloctyl]thiophene-2-sulfonamide
58	20	10	5-bromo-N-[(1S)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide
59	20	4.5	5-bromo-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide
60	20	6.9	5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide
61	20	5.7	5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-4-methyl-2-propylpentyl]thiophene-2-sulfonamide
62	20	4.5	5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)heptyl]thiophene-2-sulfonamide
63	20	3.3	5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-isobutylheptyl]thiophene-2-sulfonamide
64	20	2.9	5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)heptyl]thiophene-2-sulfonamide
65	20	1.9	5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-pentyloctyl]thiophene-2-sulfonamide
66	20	5.3	5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide
67	20	4.7	5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-phenylbutyl]thiophene-2-sulfonamide
68	20	2.3	5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-4-methyl-2-phenylpentyl]thiophene-2-sulfonamide
69	20	1.5	5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-4-methyl-2-pyridin-3-ylpentyl]thiophene-2-sulfonamide
70	20	4.6	5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide
71	20	5.2	5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide
72	20	6.4	5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide

73	20	1.9	5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide
74	20	5.1	5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)-3-methylbutyl]thiophene-2-sulfonamide
75	20	4.7	5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-isopropyl-4-methylpentyl]thiophene-2-sulfonamide
76	20	3.2	N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)butyl]-5-bromothiophene-2-sulfonamide
77A	20	3.1	5-chloro-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide
77B	20	16.9	5-chloro-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide
78A	20	5.8	5-bromo-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide
78B	20	23.0	5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide
79	20	1.6	5-chloro-N-[(1S)-1-(hydroxymethyl)-2-(methylamino)butyl]-2-thiophenesulfonamide
80	20	1.8	5-chloro-N-[(1S)-2-(ethylamino)-2-(hydroxymethyl)propyl]-2-thiophenesulfonamide
81	20	1.5	5-chloro-N-[(1S)-2-[(2-hydroxyethyl)amino]-1-(hydroxymethyl)propyl]-2-thiophenesulfonamide
82	20	2.2	5-chloro-N-[(1S)-2-[(2-hydroxyethyl)amino]-1-(hydroxymethyl)butyl]-2-thiophenesulfonamide
83	20	1.7	5-chloro-N-[(1S)-2-[(2-hydroxyethyl)amino]-1-(hydroxymethyl)heptyl]-2-thiophenesulfonamide
84	20	3.8	N-[(1S)-2-(benzylamino)-1-(hydroxymethyl)propyl]-5-chloro-2-thiophenesulfonamide
85	20	8.9	N-[(1S)-2-(benzylamino)-1-(hydroxymethyl)butyl]-5-chloro-2-thiophenesulfonamide
86	20	20	5-chloro-N-[(1S)-2-(cyclopentylamino)-1-(hydroxymethyl)propyl]-2-thiophenesulfonamide
87	20	3.1	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-N-(2-phenoxyethyl)thiophene-2-sulfonamide
88	20	4.9	5-chloro-N-(3-chlorobenzyl)-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
89	20	2.3	5-chloro-N-[(S)-2-hydroxy-1-phenylethyl]thiophene-2-sulfonamide
90	10 20 3	3.82 3.6 1.74	5-chloro-N-[(S)-1-(hydroxymethyl)-3-methylbutyl]thiophene-2-sulfonamide

91	20	2	5-chloro-N-[1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide
92	20	1.5	5-chloro-N-(2-hydroxy-1,1-dimethylethyl)thiophene-2-sulfonamide
93	20	2	N-[1,1-bis(hydroxymethyl)propyl]-5-chlorothiophene-2-sulfonamide
94	10	2.4	5-chloro-N-[1-(hydroxymethyl)cyclopentyl]thiophene-2-sulfonamide
	20	2	
	20	3.4	
	20	2.2	
	20	2.2	
95	20	5.8	
	10	1.9	5-chloro-N-[(S)-2-cyclohexyl-1-(hydroxymethyl)ethyl]thiophene-2-sulfonamide
	20	4.1	
	20	3.8	
	20	1.9	
96	20	1.5	N-[(S)-1-benzyl-2-hydroxyethyl]-5-chlorothiophene-2-sulfonamide
97	20	1.8	5-chloro-N-[1-(hydroxymethyl)butyl]thiophene-2-sulfonamide
98	10	2.43	5-chloro-N-[(S)-1-(hydroxymethyl)-2,2-dimethylpropyl]thiophene-2-sulfonamide
	20	2.8	
	20	4.9	
	20	2.4	
99	20	1.5	5-chloro-N-[(R,R)-2-hydroxy-1-(hydroxymethyl)-2-(4-nitrophenyl)ethyl]thiophene-2-sulfonamide
100	20	1.5	5-chloro-N-[(S)-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide
101	20	2.7	N-[R-2-(benzylthio)-1-(hydroxymethyl)ethyl]-5-chlorothiophene-2-sulfonamide
102	20	1.6	N-[(R,S)-2-(benzyloxy)-1-(hydroxymethyl)propyl]-5-chlorothiophene-2-sulfonamide
103	20	2.3	5-chloro-N-[(R,R)-2-hydroxy-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide
104	20	2.5	5-bromo-N-[(S)-2-hydroxy-1-phenylethyl]thiophene-2-sulfonamide
105	10	3.93	5-bromo-N-[(S)-1-(hydroxymethyl)-3-methylbutyl]thiophene-2-sulfonamide
	3	2.23	
	1	1.57	
	20	2.4	
	20	1.5	
106	20	2.4	5-bromo-N-[1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide
107	20	1.5	5-bromo-N-(2-hydroxy-1,1-dimethylethyl)thiophene-2-sulfonamide

108	20	2	N-[1,1-bis(hydroxymethyl)propyl]-5-bromothiophene-2-sulfonamide
109	10	2	5-bromo-N-[1-(hydroxymethyl)cyclopentyl]thiophene-2-sulfonamide
	3	1.8	
	20	2.5	
	20	1.7	
	20	2.7	
	20	4.3	
110	20	4.8	
	10	1.71	5-bromo-N-[(S)-2-cyclohexyl-1-(hydroxymethyl)ethyl]thiophene-2-sulfonamide
	20	2.9	
	20	2.3	
	20	3.6	
111	20	1.8	5-bromo-N-[(S)-1-(hydroxymethyl)-3-(methylthio)propyl]thiophene-2-sulfonamide
112	20	1.6	5-bromo-N-[1-(hydroxymethyl)butyl]thiophene-2-sulfonamide
113	10	2.86	5-bromo-N-[(S)-1-(hydroxymethyl)-2,2-dimethylpropyl]thiophene-2-sulfonamide
	20	3.7	
	20	1.7	
	20	3	
114	20	1.8	N-[R-2-(benzylthio)-1-(hydroxymethyl)ethyl]-5-bromothiophene-2-sulfonamide
115	20	4.4	5-bromo-N-(R-2-hydroxy-1-[(3-methylbenzyl)thio]methyl)ethyl]thiophene-2-sulfonamide
116	20	2.2	N-[(S)-1-[4-(benzyloxy)benzyl]-2-hydroxyethyl]-5-bromothiophene-2-sulfonamide
117	20	2.6	5-bromo-N-[(R,R)-2-hydroxy-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide
118	20	4.8	5-chloro-N-[(S,S)-1-formyl-2-methylbutyl]thiophene-2-sulfonamide
119	10	4.1	5-chloro-N-[(S,S)-1-(1-hydroxyethyl)-2-methylbutyl]thiophene-2-sulfonamide
	0.3	1.54	
	3	2.49	
	10	3.76	
120	10	8.83	5-chloro-N-[(S,S)-1-[cyclopentyl(hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide
	10	5.24	
	3	1.9	
121	10	1.74	5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]octyl]thiophene-2-sulfonamide
	10	1.5	
	10	1.56	
122	10	1.68	5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]heptyl]thiophene-2-sulfonamide
	10	1.8	
123	10	2.14	5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]hexyl]thiophene-2-sulfonamide
	10	1.76	

124	10	2.32	5-chloro-N-[(S)-2-hydroxy-3-methyl-1-[(S)-1-methylpropyl]butyl]thiophene-2-sulfonamide
	10	2.42	
125	10	6.52	5-chloro-N-[(S)-2-hydroxy-3,3-dimethyl-1-[(S)-1-methylpropyl]butyl]thiophene-2-sulfonamide
	3	1.58	
	10	3.94	
126	10	5.25	5-chloro-N-[(S)-2-hydroxy-4-methyl-1-[(S)-1-methylpropyl]pentyl]thiophene-2-sulfonamide
	10	3.31	
127	10	4.27	5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]but-3-enyl]thiophene-2-sulfonamide
128	10	5.62	5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl]thiophene-2-sulfonamide
	1	1.52	
	20	5.9	
	3	2.48	
	10	3.86	
129	10	7.43	5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]butyl]thiophene-2-sulfonamide
	3	1.61	
	10	4.59	
130	10	1.85	5-chloro-N-[(S,S)-1-[(4-fluorophenyl)(hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide
131	10	3.64	5-chloro-N-[(S,S)-1-[(4-chlorophenyl)(hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide
132	10	1.68	5-chloro-N-[(S)-2-hydroxy-4-methyl-1-[(S)-1-methylpropyl]pent-3-enyl]thiophene-2-sulfonamide
133	10	2.51	5-chloro-N-[(S)-2-hydroxy-3-methyl-1-[(S)-1-methylpropyl]but-3-enyl]thiophene-2-sulfonamide
		1.68	
134	10	2.42	5-chloro-N-[(S,S)-1-[hydroxy(4-methoxyphenyl)methyl]-2-methylbutyl]thiophene-2-sulfonamide
135	10	1.93	5-chloro-N-[(S,E)-2-hydroxy-3-methyl-1-[(S)-1-methylpropyl]pent-3-enyl]thiophene-2-sulfonamide
	10	1.75	
136	10	2.13	5-chloro-N-[(S)-4-(1,3-dioxan-2-yl)-2-hydroxy-1-[(S)-1-methylpropyl]butyl]thiophene-2-sulfonamide
137	10	1.96	5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]hex-5-enyl]thiophene-2-sulfonamide
138	10	3.54	5-chloro-N-[(S,S)-1-{hydroxy[4-(methylthio)phenyl]methyl}-2-methylbutyl]thiophene-2-sulfonamide
139	10	3.97	5-chloro-N-[(S,S)-1-[[4-(dimethylamino)phenyl](hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide
140	10	2.57	N-[(S,S)-1-[cyclopentyl(hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide
141	10	4.2	N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]octyl]thiophene-2-sulfonamide

142	10	3.59	N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]heptyl)thiophene-2-sulfonamide
143	10	1.64	N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]hexyl)thiophene-2-sulfonamide
144	10	1.51	N-((S,S)-1-[hydroxy(2-methylphenyl)methyl]-2-methylbutyl)thiophene-2-sulfonamide
145	10	1.72	N-((S)-2-hydroxy-3,3-dimethyl-1-[(S)-1-methylpropyl]butyl)thiophene-2-sulfonamide
146	10	1.83	N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]but-3-enyl)thiophene-2-sulfonamide
147	10	2.04	N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl)thiophene-2-sulfonamide
148	10	1.52	N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]butyl)thiophene-2-sulfonamide
149	10	1.62	N-((S,S)-1-[hydroxy(4-methoxyphenyl)methyl]-2-methylbutyl)thiophene-2-sulfonamide
150	10	1.6	N-((S)-4-(1,3-dioxan-2-yl)-2-hydroxy-1-[(S)-1-methylpropyl]butyl)thiophene-2-sulfonamide
151	10	1.94	N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]hex-5-enyl)thiophene-2-sulfonamide
152	10	1.51	N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-3-ynyl)thiophene-2-sulfonamide
153	10	2.09	N-((S,S)-1-{hydroxy[4-(methylthio)phenyl]methyl}-2-methylbutyl)thiophene-2-sulfonamide
154	10	4.23	N-((S,S)-1-[[4-(dimethylamino)phenyl](hydroxy)methyl]-2-methylbutyl)thiophene-2-sulfonamide
155	20 20	4.2 2.6	5-chloro-N-((S,S)-1-[(S)-cyclohex-2-en-1-yl(hydroxy)methyl]-2-methylbutyl)thiophene-2-sulfonamide
156	20 20	5.3 5.9	5-chloro-N-((S,S,E)-2-hydroxy-1-[(S)-1-methylpropyl]hex-4-enyl)thiophene-2-sulfonamide
157	20 20	6.7 6.8	5-chloro-N-((S,R,E)-2-hydroxy-1-[(S)-1-methylpropyl]hex-4-enyl)thiophene-2-sulfonamide
158	20 20	1.6 5.4	5-chloro-N-((S,R,E)-2-hydroxy-1-[(S)-1-methylpropyl]hept-4-enyl)thiophene-2-sulfonamide
159	20 20	3.8 3.2	5-chloro-N-((S,S)-2-hydroxy-4-methyl-1-[(S)-1-methylpropyl]pent-4-enyl)thiophene-2-sulfonamide
160	20 20	4 3.3	5-chloro-N-((S,R)-2-hydroxy-4-methyl-1-[(S)-1-methylpropyl]pent-4-enyl)thiophene-2-sulfonamide
161	20	2.6	5-chloro-N-((S,E)-2-hydroxy-1-[(S)-1-methylpropyl]-5-phenylpent-4-enyl)thiophene-2-sulfonamide
162	10	2.6	5-chloro-N-[(S,S)-1-(1-hydroxy-1-methylethyl)-2-methylbutyl]thiophene-2-sulfonamide
163	10	3.3	5-chloro-N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]-2-pentylheptyl)thiophene-2-sulfonamide

164	10	1.72	5-chloro-N-((S,S)-1-[hydroxy(diphenyl)methyl]-2-methylbutyl)thiophene-2-sulfonamide
165	10	1.73	N-((S)-2-allyl-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl)-5-chlorothiophene-2-sulfonamide
166	10	1.78	5-chloro-N-((S)-2-ethyl-2-hydroxy-1-[(S)-1-methylpropyl]butyl)thiophene-2-sulfonamide
167	10	3.42	N-((S,S)-1-[bis(4-chlorophenyl)(hydroxy)methyl]-2-methylbutyl)-5-chlorothiophene-2-sulfonamide
168	10	5.87	5-chloro-N-((S)-2-hydroxy-2-isopropenyl-3-methyl-1-[(S)-1-methylpropyl]but-3-enyl)thiophene-2-sulfonamide
	10	1.6	
	3	1.5	
169	10	1.51	5-chloro-N-((S,S)-1-{hydroxy[bis(4-methoxyphenyl)]methyl}-2-methylbutyl)thiophene-2-sulfonamide
170	10	1.95	5-chloro-N-((S,E)-2-hydroxy-3-methyl-2-[(E)-1-methylprop-1-enyl]-1-[(S)-1-methylpropyl]pent-3-enyl)thiophene-2-sulfonamide
171	10	5.32	N-((S)-2-but-3-enyl-2-hydroxy-1-[(S)-1-methylpropyl]hex-5-enyl)-5-chlorothiophene-2-sulfonamide
172	10	2	5-chloro-N-((S,S)-1-{hydroxy[di(1-naphthyl)]methyl}-2-methylbutyl)thiophene-2-sulfonamide
173	10	1.5	5-bromo-N-((S)-2-ethyl-2-hydroxy-1-[(S)-1-methylpropyl]butyl)thiophene-2-sulfonamide
174	10	1.75	5-bromo-N-((S)-2-hydroxy-2-isopropenyl-3-methyl-1-[(S)-1-methylpropyl]but-3-enyl)thiophene-2-sulfonamide
175	10	2.27	5-bromo-N-((S,E)-2-hydroxy-3-methyl-2-[(E)-1-methylprop-1-enyl]-1-[(S)-1-methylpropyl]pent-3-enyl)thiophene-2-sulfonamide
	10	1.7	
	20	3.3	
176	10	1.52	5-bromo-N-((S)-2-but-3-enyl-2-hydroxy-1-[(S)-1-methylpropyl]hex-5-enyl)thiophene-2-sulfonamide
177	20	5.1	5-chloro-N-[1-(hydroxymethyl)cyclohexyl]thiophene-2-sulfonamide
	20	3.7	
178	20	2.3	5-chloro-N-[2-(hydroxymethyl)bicyclo[2.2.1]hept-2-yl]thiophene-2-sulfonamide
179	20	8.3	5-chloro-N-[1-(hydroxymethyl)-2,3-dihydro-H-inden-1-yl]thiophene-2-sulfonamide
180	20	2.3	5-chloro-N-[2-(hydroxymethyl)-2,3-dihydro-H-inden-2-yl]thiophene-2-sulfonamide
181	20	2.6	5-bromo-N-[1-(hydroxymethyl)cyclohexyl]thiophene-2-sulfonamide
182	20	4.2	5-bromo-N-[2-(hydroxymethyl)bicyclo[2.2.1]hept-2-yl]thiophene-2-sulfonamide
183	20	3.7	5-bromo-N-[2-(hydroxymethyl)-2,3-dihydro-H-inden-2-yl]thiophene-2-sulfonamide

184	20	3.7	5-chloro-N-((S,S)-1-[(S)-1-hydroxyethyl]-2-methylbutyl)thiophene-2-sulfonamide
	20	2.7	
185	20	2.4	5-chloro-N-((S,S)-1-[R-1-hydroxyethyl]-2-methylbutyl)thiophene-2-sulfonamide
	20	5.4	
186	20	6.3	5-chloro-N-((S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl)thiophene-2-sulfonamide
	20	2.4	
187	20	5.7	5-chloro-N-((S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl)thiophene-2-sulfonamide
	20	3	
188	20	6.3	5-chloro-N-((S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl)thiophene-2-sulfonamide
	20	3.6	
	20	3.6	
189	20	8.1	5-chloro-N-((S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl)thiophene-2-sulfonamide
	20	7.9	
	20	7.2	
	20	4.4	
190	20	4.4	5-bromo-N-((S,S)-1-[(S)-1-hydroxyethyl]-2-methylbutyl)thiophene-2-sulfonamide
	20	2.6	
191	20	5.4	5-bromo-N-((S,S)-1-[R-1-hydroxyethyl]-2-methylbutyl)thiophene-2-sulfonamide
	20	4.8	
192	20	2.2	5-bromo-N-((S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl)thiophene-2-sulfonamide
	20	1.9	
	20	1.7	
193	20	8.5	5-bromo-N-((S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl)thiophene-2-sulfonamide
	20	4.5	
	20	3	
194	20	9	5-bromo-N-((S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl)thiophene-2-sulfonamide
	20	5.3	
195	20	4.2	5-bromo-N-((S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl)thiophene-2-sulfonamide
	20	6.7	
196	20	20.7	5-chloro-N-[(S,S)-2-methyl-1-(2,2,2-trifluoro-1-hydroxyethyl)butyl]thiophene-2-sulfonamide
197	20	3.4	5-chloro-N-[1-(1-hydroxybut-3-enyl)cyclohexyl]thiophene-2-sulfonamide
	20	1.8	
198	20	4.4	5-chloro-N-[1-(1-hydroxy-3-methylbut-3-enyl)cyclohexyl]thiophene-2-sulfonamide
199	20	2.8	5-chloro-N-[(S)-2-hydroxy-1-(4-methoxycyclohexyl)ethyl]thiophene-2-sulfonamide
199A	20	2.4	5-Chloro-N-[(S)-2-hydroxy-1-(4-hydroxycyclohexyl)ethyl]thiophene-2-sulfonamide
200	20	2	5-chloro-N-[(S)-2-hydroxy-1-(4-propoxycyclohexyl)ethyl]thiophene-2-sulfonamide
201	20	2.2	N-[(S)-1-[4-(allyloxy)cyclohexyl]-2-hydroxyethyl]-5-chlorothiophene-2-sulfonamide
202	20	2	N-[(S)-1-[4-(benzyloxy)cyclohexyl]-2-hydroxyethyl]-5-chlorothiophene-2-sulfonamide

203	20	1.5	N-[1-acetyl-4-(hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide
204	20 20	2.8 2.5	5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide
205	20	2.6	N-[(1S)-2-butyl-1-(hydroxymethyl)hexyl]-5-chloro-2-thiophenesulfonamide
206	20	1.8	N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide
207	20	5.6	N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-5-iodo-2-thiophenesulfonamide
208	20	16.5	5-fluoro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide
209	20	2.1	4-[1-(5-chloro-thiophene-2-sulfonylamino)-2-hydroxyethyl]-piperidine-1-carboxylic acid tert-butyl ester
210	20	7.7	N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide
211	20	1.9	5-chloro-N-[(S)-2-hydroxy-1-(4-benzylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
212	20	1.7	5-chloro-N-[(S)-2-hydroxy-1-(4-methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
213	20	1.6	5-chloro-N-[(S)-2-hydroxy-1-(4-ethylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
214	20	2.1	5-chloro-N-[(S)-2-hydroxy-1-(4-npropylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
215	20	2.0	5-chloro-N-[(S)-2-hydroxy-1-(4-allylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
216	20	1.5	5-chloro-N-[(S)-2-hydroxy-1-(4-(3-pyridyl)methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
217	20	1.8	5-chloro-N-[(S)-2-hydroxy-1-(4-morpholinocyclohexyl)ethyl]thiophene-2-sulfonamide
218	20	2.8	5-chloro-N-[(S)-2-hydroxy-1-(4-(4-pyridyl)methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
219	20	2.3	5-chloro-N-[(S)-2-hydroxy-1-(4-(2-pyridyl)methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide
220	20	2.5	5-chloro-N-[(S)-2-hydroxy-1-(4-(carboethoxymethyl)aminocyclohexyl)ethyl]thiophene-2-sulfonamide
221	20	4.9	5-chloro-N-[(S)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide


222	20	7.2 9.7	5-chloro- <i>N</i> -[(<i>S</i>)-2-ethyl-1-(1-hydroxyethyl)butyl]thiophene-2-sulfonamide
223	20	5.6 8.4	5-chloro- <i>N</i> -[(<i>S</i>)-2-ethyl-1-(1-hydroxy-1-methylethyl)butyl]thiophene-2-sulfonamide
224	20	7.2	5-chloro- <i>N</i> -(2-hydroxy-1-tetrahydro- <i>H</i> -thiopyran-4-ylethyl)thiophene-2-sulfonamide
225	20	8.0	5-chloro- <i>N</i> -[(<i>S</i>)-2-hydroxy-1-piperidin-4-ylethyl]thiophene-2-sulfonamide
226	20	7.6	<i>N</i> -[(<i>S</i>)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide
227	20	26.3	<i>N</i> -[(<i>S</i>)-2-ethyl-1-(hydroxymethyl)butyl]-5-fluorothiophene-2-sulfonamide
228	20	9.9	5-chloro- <i>N</i> -[(<i>S</i>)-1-(2,3-dihydro- <i>H</i> -inden-2-yl)-2-hydroxyethyl]thiophene-2-sulfonamide
229	20	6.3	5-chloro- <i>N</i> -{(<i>S,S</i>)-1-[(<i>Z</i>)-(hydroxyimino)methyl]-2-methylbutyl}thiophene-2-sulfonamide
230	20	4.8	5-chloro- <i>N</i> -{(<i>S,S</i>)-1-[(<i>E</i>)-(hydroxyimino)methyl]-2-methylbutyl}thiophene-2-sulfonamide

All publications cited in this specification are incorporated herein by reference. While the invention has been described with reference to a particularly preferred embodiment, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

What is claimed is:

1. A compound of Formula (I), or pharmaceutically acceptable salt thereof, wherein Formula (I) has the structure:

5



(I)

wherein:

10 R₁ and R₂ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, CF₃, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, phenyl, substituted phenyl, and (CH₂)_n(1,3)dioxane, where n is 2 to 5;

15 R₃ is selected from the group consisting of hydrogen, alkyl, and substituted alkyl;

R₄ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkylcycloalkyl, substituted alkylcycloalkyl, phenyl(substituted)alkyl, alkylOH, substituted alkylOH, alkylOBn, substituted alkylOBn, alkylpyridyl, substituted alkylpyridyl, alkylfuranyl, substituted alkylfuranyl, CH(OH)phenyl, CH(OH)substituted phenyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, N-substituted-piperidiny, piperidiny, substituted piperidiny, tetrahydrothiopyran, substituted tetrahydrothiopyran, 2-indane, substituted 2-indane, phenyl, substituted phenyl, alkylNHR₇, and substituted alkylNHR₇;

with the proviso that R₃ and R₄ are not both hydrogen;

25 R₇ is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, benzyl, substituted benzyl, alkylOH, substituted alkylOH, alkylSR₈, or substituted alkylSR₈;

R₈ is alkyl, substituted alkyl, benzyl, or substituted benzyl;

or R₃ and R₄ may be joined to form a ring;


R_3 is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, $\text{CH}_2\text{cycloalkyl}$, substituted $\text{CH}_2\text{cycloalkyl}$, benzyl, substituted benzyl, and $\text{CH}_2\text{CH}_2\text{QR}_9$;

Q is O, NH or S;

5 R_9 is lower alkyl, substituted lower alkyl, phenyl, or substituted phenyl;

R_6 is selected from the group consisting of hydrogen, halogen and CF_3 ;

T is selected from the group consisting of



10 W, Y and Z are independently selected from the group consisting of C, CR_{10} and N;

R_{10} is selected from the group consisting of hydrogen and halogen, with the proviso that at least one of W, Y and Z must be C;


X is selected from the group consisting of O, S, SO_2 , and NR_{11} ;

15 R_{11} is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, and substituted phenyl;

provided that when the compound contains one or more chiral centers, at least one of the chiral centers must be of S-stereochemistry.

2. A compound of Formula (I), or pharmaceutically acceptable salt thereof, wherein Formula (I) has the structure:

20



(I)

wherein:

R_1 and R_2 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, CF_3 , alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, phenyl, substituted phenyl, and

5 $(CH_2)_n(1,3)$ dioxane, where n is 2 to 5;

R_3 is selected from the group consisting of hydrogen, alkyl, and substituted alkyl;

R_4 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkylcycloalkyl, substituted alkylcycloalkyl, phenyl(substituted)alkyl, alkylOH, substituted alkylOH, alkylOBn, substituted alkylOBn, alkylpyridyl, substituted alkylpyridyl, alkylfuranyl, substituted alkylfuranyl, $CH(OH)$ phenyl, $CH(OH)$ substituted phenyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, N-substituted-piperidiny, piperidiny, substituted piperidiny, tetrahydrothiopyran, substituted tetrahydrothiopyran, 2-indane, substituted 2-indane, phenyl, substituted phenyl, alkylNHR₇, and substituted alkylNHR₇;

with the proviso that R_3 and R_4 are not both hydrogen;

R_7 is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, benzyl, substituted benzyl, alkylOH, substituted alkylOH, alkylSR₈, or substituted alkylSR₈;

R_8 is alkyl, substituted alkyl, benzyl, or substituted benzyl;

20 or R_3 and R_4 may be joined to form a ring;


R_5 is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CH_2 cycloalkyl, substituted CH_2 cycloalkyl, benzyl, substituted benzyl, and $CH_2CH_2QR_9$;

Q is O, NH or S;

25 R_9 is lower alkyl, substituted lower alkyl, phenyl, or substituted phenyl;

R_6 is selected from the group consisting of hydrogen, halogen and CF_3 ;

T is



W is C, wherein W provides the point of attachment to the SO₂ group;


X is S;

Y and Z are independently C or CR₁₀;

R₁₀ is selected from the group consisting of hydrogen and halogen;

- 5 provided that when the compound contains one or more chiral centers, at least one of the chiral centers must be of S-stereochemistry.

3. A compound of Formula (I), or pharmaceutically acceptable salt thereof, wherein Formula (I) has the structure:



10

(I)

wherein:

- 15 R₁ and R₂ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, CF₃, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, phenyl, substituted phenyl, and (CH₂)_n(1,3)dioxane, where n is 2 to 5;

R₃ is selected from the group consisting of hydrogen, alkyl, and substituted alkyl;

- 20 R₄ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkylcycloalkyl, substituted alkylcycloalkyl, phenyl(substituted)alkyl, alkylOH, substituted alkylOH, alkylOBn, substituted alkylOBn, alkylpyridyl, substituted alkylpyridyl, alkylfuranyl, substituted alkylfuranyl, CH(OH)phenyl, CH(OH)substituted phenyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, N-substituted-piperidiny, piperidiny, substituted piperidiny,
- 25 tetrahydrothiopyran, substituted tetrahydrothiopyran, 2-indane, substituted 2-indane, phenyl, substituted phenyl, alkylNHR₇, and substituted alkylNHR₇;

with the proviso that R₃ and R₄ are not both hydrogen;

R_7 is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, benzyl, substituted benzyl, alkylOH, substituted alkylOH, alkylSR₈, or substituted alkylSR₈;

R_8 is alkyl, substituted alkyl, benzyl, or substituted benzyl;

or R_3 and R_4 may be joined to form a ring;


- 5 R_5 is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CH₂cycloalkyl, substituted CH₂cycloalkyl, benzyl, substituted benzyl, and CH₂CH₂QR₉;

Q is O, NH or S;

R_9 is lower alkyl, substituted lower alkyl, phenyl, or substituted phenyl;

- 10 R_6 is halogen;

T is



W is C, wherein W provides the point of attachment to the SO₂ group;


X is S;

- 15 Y and Z are independently C or CR₁₀;

R_{10} is selected from the group consisting of hydrogen and halogen;

provided that when the compound contains one or more chiral centers, at least one of the chiral centers must be of S-stereochemistry.


- 20 4. A compound of Formula (I), or pharmaceutically acceptable salt thereof, wherein Formula (I) has the structure:



(I)


- 25 wherein:

R_1 and R_2 are hydrogen;
 R_3 is hydrogen;
 R_4 is a lower alkyl of S-stereochemistry;
 R_5 is hydrogen;
 R_6 is halogen;
 T is



X is S;
 W, Y and Z are C, wherein W provides the point of attachment to the SO₂
 group.


5. A compound of Formula (I), or pharmaceutically acceptable salt thereof, wherein Formula (I) has the structure:



(I)

wherein:

R_1 and R_2 are hydrogen;
 R_3 is hydrogen;
 R_4 is a lower alkyl of S-stereochemistry;
 R_5 is hydrogen;
 R_6 is hydrogen or halogen;
 T is




X is O;

W, Y and Z are C, wherein W provides the point of attachment to the SO₂ group.

5

6. A compound of Formula (I), or pharmaceutically acceptable salt thereof, wherein Formula (I) has the structure:



(I)

10 wherein:

R₁ and R₂ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, CF₃, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, phenyl, substituted phenyl, and (CH₂)_n(1,3)dioxane, where n is 2 to 5;

15 R₃ is selected from the group consisting of hydrogen, alkyl, and substituted alkyl;

R₄ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkylcycloalkyl, substituted alkylcycloalkyl, phenyl(substituted)alkyl, alkylOH, substituted alkylOH, alkylOBn, substituted alkylOBn, alkylpyridyl, substituted alkylpyridyl, alkylfuranyl, substituted alkylfuranyl, CH(OH)phenyl, CH(OH)substituted phenyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, N-substituted-piperidiny, piperidiny, substituted piperidiny, tetrahydrothiopyran, substituted tetrahydrothiopyran, 2-indane, substituted 2-indane, phenyl, substituted phenyl, alkylNHR₇, and substituted alkylNHR₇;

25 with the proviso that R₃ and R₄ are not both hydrogen;

R₇ is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, benzyl, substituted benzyl, alkylOH, substituted alkylOH, alkylSR₈, or substituted alkylSR₈;

R₈ is alkyl, substituted alkyl, benzyl, or substituted benzyl;

or R_3 and R_4 may be joined to form a ring;


R_5 is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, $\text{CH}_2\text{cycloalkyl}$, substituted $\text{CH}_2\text{cycloalkyl}$, benzyl, substituted benzyl, and $\text{CH}_2\text{CH}_2\text{QR}_9$;

5 Q is O, NH or S;

R_9 is lower alkyl, substituted lower alkyl, phenyl, or substituted phenyl;

R_6 is selected from the group consisting of hydrogen, halogen and CF_3 ;

T is



10 W , Y and Z are independently selected from the group consisting of C, CR_{10} and N;


R_{10} is selected from the group consisting of hydrogen and halogen, with the proviso that at least one of W , Y and Z must be C;

X is selected from the group consisting of O, S, SO_2 , and NR_{11} ;

15 R_{11} is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, and substituted phenyl;

provided that when the compound contains one or more chiral centers, at least one of the chiral centers must be of S-stereochemistry.

20 7. A compound of Formula (I), or pharmaceutically acceptable salt thereof, wherein Formula (I) has the structure:



(I)

wherein:

R_1 and R_2 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, CF_3 , alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, phenyl, substituted phenyl, and $(CH_2)_n(1,3)$ dioxane, where n is 2 to 5;

5 R_3 is selected from the group consisting of hydrogen, alkyl, and substituted alkyl;

R_4 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkylcycloalkyl, substituted alkylcycloalkyl, phenyl(substituted)alkyl, alkylOH, substituted alkylOH, alkylOBn, substituted alkylOBn, alkylpyridyl, substituted
10 alkylpyridyl, alkylfuranyl, substituted alkylfuranyl, $CH(OH)$ phenyl, $CH(OH)$ substituted phenyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, N-substituted-piperidiny, piperidiny, substituted piperidiny, tetrahydrothiopyran, substituted tetrahydrothiopyran, 2-indane, substituted 2-indane, phenyl, substituted phenyl, alkylNHR₇, and substituted alkylNHR₇;

15 with the proviso that R_3 and R_4 are not both hydrogen;

R_7 is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, benzyl, substituted benzyl, alkylOH, substituted alkylOH, alkylSR₈, or substituted alkylSR₈;

R_8 is alkyl, substituted alkyl, benzyl, or substituted benzyl;

or R_3 and R_4 may be joined to form a ring;


20 R_5 is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, CH_2 cycloalkyl, benzyl, substituted benzyl, and $CH_2CH_2QR_9$;

Q is O, NH or S;

R_9 is lower alkyl, substituted lower alkyl, phenyl, or substituted phenyl;

25 R_6 is selected from the group consisting of hydrogen, halogen and CF_3 ;

T is



W , Y and Z are independently selected from the group consisting of C, CR₁₀ and N;

R_{10} is selected from the group consisting of hydrogen and halogen, with the proviso that at least one of W, Y and Z must be C;

X is selected from the group consisting of O, S, SO_2 , and NR_{11} ;

R_{11} is selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, and substituted phenyl;
5 provided that when the compound contains one or more chiral centers, at least one of the chiral centers must be of S-stereochemistry.

8. The compound according to any of claims 1, 2, or 5 to 7, wherein R_6 is
10 halogen.

9. The compound according to any of claims 1 to 8, wherein R_6 is chlorine or bromine.

15 10. The compound according to any of claims 1 to 3, wherein T is $C(OH)R_1R_2$ and R_1 and R_2 are each hydrogen.

11. The compound according to any of claims 1 to 7, wherein W and Z are both C.
20

12. The compound according to any of claims 1 to 3, 6 or 7, wherein R_4 is lower alkyl of S-stereochemistry.

13. The compound according to any of claims 1 to 3 or 5 to 7, wherein X
25 is S, W is C, Z is C, R_6 is halogen, R_4 is lower alkyl of S-stereochemistry, R_3 is hydrogen, R_5 is hydrogen, and R_1 and R_2 are each hydrogen.

14. The compound according to any of claims 1 to 3, 6, or 7, wherein R_3CR_4 is cyclohexyl.
30

15. The compound according to any of claims 1 to 3, 6, or 7, wherein R_3CR_4 is piperidine or N-substituted piperidine.

16. The compound according to any of claims 1 to 3, 6, or 7, wherein X is S, and W, Y and Z are independently C or CR₁₀.

5 17. The compound according to claim 1, wherein the compound is selected from the group consisting of:

3-bromo-5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;

10 5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;

4-bromo-5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;

5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;

15 2,5-dichloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-3-sulfonamide;

4,5-dichloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;

20 N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;

5-chloro-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide;

25 5-bromo-N-[(1R)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide;

4,5-dibromo-N-[(1S)-1-(hydroxymethyl)-2-methylpropyl]thiophene-2-sulfonamide;

30 5-chloro-N-[(1S)-1-cyclohexyl-2-hydroxyethyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S)-1-cyclohexyl-2-hydroxyethyl]thiophene-2-sulfonamide;

- 5-chloro-N-[1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide;
- 5-bromo-N-[1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide;
- 5 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;
- 5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;
- 5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide 1,1-dioxide;
- 10 5-chloro-N-[1-(hydroxymethyl)-2,3-dimethylpentyl]thiophene-2-sulfonamide;
- 5-chloro-N-[1-(hydroxymethyl)-2-methylpentyl]thiophene-2-sulfonamide;
- 15 5-chloro-N-[2-ethyl-1-(hydroxymethyl)hexyl]thiophene-2-sulfonamide;
- 5-chloro-N-[2-hydroxy-1-(2,4,6-trimethylcyclohex-3-en-1-yl)ethyl]thiophene-2-sulfonamide;
- 5-chloro-N-(1-cyclohex-3-en-1-yl-2-hydroxyethyl)thiophene-2-sulfonamide;
- 20 5-chloro-N-(1-cyclopentyl-2-hydroxyethyl)thiophene-2-sulfonamide;
- 5-bromo-N-[(1S)-1-(hydroxymethyl)-1,2-dimethylpropyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S)-1-(hydroxymethyl)-1,2-dimethylpropyl]thiophene-2-sulfonamide;
- 25 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2,4-dimethylpentyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)propyl]thiophene-2-sulfonamide;
- 30 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-methyloctyl]thiophene-2-sulfonamide;

- 5-chloro-N-[(1S)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide;
- 5 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)butyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylpentyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide;
- 10 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-4-methyl-2-propylpentyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)pentyl]thiophene-2-sulfonamide;
- 15 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-propyloctyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-phenylpentyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylheptyl]thiophene-2-sulfonamide;
- 20 5-chloro-N-[(1S)-2-propyl-1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)heptyl]thiophene-2-sulfonamide;
- 25 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-isobutylheptyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)heptyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-pentyloctyl]thiophene-2-sulfonamide;
- 30 5-chloro-N-[(1S,2R)-1-(hydroxymethyl)-2-phenylheptyl]thiophene-2-sulfonamide;

- 5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-4-methyl-2-phenylpentyl]thiophene-2-sulfonamide;
- 5 5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide;
- 10 5-chloro-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-isopropyloctyl]thiophene-2-sulfonamide;
- 15 N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)propyl]-5-chlorothiophene-2-sulfonamide;
- N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)butyl]-5-chlorothiophene-2-sulfonamide;
- N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)-4-methylpentyl]-5-chlorothiophene-2-sulfonamide;
- 20 5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2,4-dimethylpentyl]thiophene-2-sulfonamide;
- 5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-methyloctyl]thiophene-2-sulfonamide;
- 25 5-bromo-N-[(1S)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide;
- 5-bromo-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide;
- 5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide;
- 30 5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-4-methyl-2-propylpentyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)heptyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-isobutylheptyl]thiophene-2-sulfonamide;

5 5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-(4-methoxyphenyl)heptyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2R)-1-(hydroxymethyl)-2-pentyl-octyl]thiophene-2-sulfonamide;

10 5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-phenylpropyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-phenylbutyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-4-methyl-2-phenylpentyl]thiophene-2-sulfonamide;

15 5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-4-methyl-2-pyridin-3-ylpentyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide;

20 5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)-4-methylpentyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2R)-2-(2-furyl)-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide;

25 5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)-3-methylbutyl]thiophene-2-sulfonamide;

5-bromo-N-[(1S,2S)-1-(hydroxymethyl)-2-isopropyl-4-methylpentyl]thiophene-2-sulfonamide;

30 N-[(1S,2S)-2-[1,1'-biphenyl]-4-yl-1-(hydroxymethyl)butyl]-5-bromothiophene-2-sulfonamide;

5-chloro-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide;

- 5-chloro-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide;
- 5-bromo-N-[(1S,2R)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide;
- 5 5-bromo-N-[(1S,2S)-2-ethyl-1-(hydroxymethyl)octyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(1S)-1-(hydroxymethyl)-2-(methylamino)butyl]-2-thiophenesulfonamide;
- 5-chloro-N-[(1S)-2-(ethylamino)-2-(hydroxymethyl)propyl]-2-thiophenesulfonamide;
- 10 thiophenesulfonamide;
- 5-chloro-N-[(1S)-2-[(2-hydroxyethyl)amino]-1-(hydroxymethyl)propyl]-2-thiophenesulfonamide;
- 5-chloro-N-[(1S)-2-[(2-hydroxyethyl)amino]-1-(hydroxymethyl)butyl]-2-thiophenesulfonamide;
- 15 5-chloro-N-[(1S)-2-[(2-hydroxyethyl)amino]-1-(hydroxymethyl)heptyl]-2-thiophenesulfonamide;
- N-[(1S)-2-(benzylamino)-1-(hydroxymethyl)propyl]-5-chloro-2-thiophenesulfonamide;
- N-[(1S)-2-(benzylamino)-1-(hydroxymethyl)butyl]-5-chloro-2-thiophenesulfonamide;
- 20 thiophenesulfonamide;
- 5-chloro-N-[(1S)-2-(cyclopentylamino)-1-(hydroxymethyl)propyl]-2-thiophenesulfonamide;
- 5-chloro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-N-(2-phenoxyethyl)thiophene-2-sulfonamide;
- 25 5-chloro-N-(3-chlorobenzyl)-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(S)-2-hydroxy-1-phenylethyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(S)-1-(hydroxymethyl)-3-methylbutyl]thiophene-2-sulfonamide;
- 30 5-chloro-N-[1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide;
- 5-chloro-N-(2-hydroxy-1,1-dimethylethyl)thiophene-2-sulfonamide;
- N-[1,1-bis(hydroxymethyl)propyl]-5-chlorothiophene-2-sulfonamide;

- 5-chloro-N-[1-(hydroxymethyl)cyclopentyl]thiophene-2-sulfonamide;
 5-chloro-N-[(S)-2-cyclohexyl-1-(hydroxymethyl)ethyl]thiophene-2-sulfonamide;
- 5 N-[(S)-1-benzyl-2-hydroxyethyl]-5-chlorothiophene-2-sulfonamide;
 5-chloro-N-[1-(hydroxymethyl)butyl]thiophene-2-sulfonamide;
 5-chloro-N-[(S)-1-(hydroxymethyl)-2,2-dimethylpropyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(R,R)-2-hydroxy-1-(hydroxymethyl)-2-(4-nitrophenyl)ethyl]thiophene-2-sulfonamide;
- 10 5-chloro-N-[(S)-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide;
 N-[R-2-(benzylthio)-1-(hydroxymethyl)ethyl]-5-chlorothiophene-2-sulfonamide;
- N-[(R,S)-2-(benzyloxy)-1-(hydroxymethyl)propyl]-5-chlorothiophene-2-sulfonamide;
- 15 5-chloro-N-[(R,R)-2-hydroxy-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide;
- 5-bromo-N-[(S)-2-hydroxy-1-phenylethyl]thiophene-2-sulfonamide;
 5-bromo-N-[(S)-1-(hydroxymethyl)-3-methylbutyl]thiophene-2-sulfonamide;
- 20 5-bromo-N-[1-(hydroxymethyl)pentyl]thiophene-2-sulfonamide;
 5-bromo-N-(2-hydroxy-1,1-dimethylethyl)thiophene-2-sulfonamide;
 N-[1,1-bis(hydroxymethyl)propyl]-5-bromothiophene-2-sulfonamide
 5-bromo-N-[1-(hydroxymethyl)cyclopentyl]thiophene-2-sulfonamide
 5-bromo-N-[(S)-2-cyclohexyl-1-(hydroxymethyl)ethyl]thiophene-2-sulfonamide;
- 25 5-bromo-N-[(S)-1-(hydroxymethyl)-3-(methylthio)propyl]thiophene-2-sulfonamide;
- 5-bromo-N-[1-(hydroxymethyl)butyl]thiophene-2-sulfonamide;
 5-bromo-N-[(S)-1-(hydroxymethyl)-2,2-dimethylpropyl]thiophene-2-sulfonamide;
- 30 N-[R-2-(benzylthio)-1-(hydroxymethyl)ethyl]-5-bromothiophene-2-sulfonamide;

- 5-bromo-N-(R-2-hydroxy-1-[(3-methylbenzyl)thio]methyl)ethylthiophene-2-sulfonamide;
N-[(S)-1-[4-(benzyloxy)benzyl]-2-hydroxyethyl]-5-bromothiophene-2-sulfonamide;
- 5 5-bromo-N-[(R,R)-2-hydroxy-1-(hydroxymethyl)propyl]thiophene-2-sulfonamide;
5-chloro-N-[(S,S)-1-formyl-2-methylbutyl]thiophene-2-sulfonamide;
5-chloro-N-[(S,S)-1-(1-hydroxyethyl)-2-methylbutyl]thiophene-2-sulfonamide;
- 10 5-chloro-N-[(S,S)-1-[cyclopentyl(hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide;
5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]octyl]thiophene-2-sulfonamide;
5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]heptyl]thiophene-2-sulfonamide;
- 15 5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]hexyl]thiophene-2-sulfonamide;
5-chloro-N-[(S)-2-hydroxy-3-methyl-1-[(S)-1-methylpropyl]butyl]thiophene-2-sulfonamide;
- 20 5-chloro-N-[(S)-2-hydroxy-3,3-dimethyl-1-[(S)-1-methylpropyl]butyl]thiophene-2-sulfonamide;
5-chloro-N-[(S)-2-hydroxy-4-methyl-1-[(S)-1-methylpropyl]pentyl]thiophene-2-sulfonamide;
5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]but-3-enyl]thiophene-2-sulfonamide;
- 25 5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl]thiophene-2-sulfonamide;
5-chloro-N-[(S)-2-hydroxy-1-[(S)-1-methylpropyl]butyl]thiophene-2-sulfonamide;
- 30 5-chloro-N-[(S,S)-1-[(4-fluorophenyl)(hydroxy)methyl]-2-methylbutyl]thiophene-2-sulfonamide;

- 5-chloro-N-((S,S)-1-((4-chlorophenyl)(hydroxy)methyl)-2-methylbutyl)thiophene-2-sulfonamide;
- 5-chloro-N-((S)-2-hydroxy-4-methyl-1-((S)-1-methylpropyl)pent-3-enyl)thiophene-2-sulfonamide;
- 5 5-chloro-N-((S)-2-hydroxy-3-methyl-1-((S)-1-methylpropyl)but-3-enyl)thiophene-2-sulfonamide;
- 5-chloro-N-((S,S)-1-[hydroxy(4-methoxyphenyl)methyl]-2-methylbutyl)thiophene-2-sulfonamide;
- 5-chloro-N-((S,E)-2-hydroxy-3-methyl-1-((S)-1-methylpropyl)pent-3-enyl)thiophene-2-sulfonamide;
- 10 5-chloro-N-((S)-4-(1,3-dioxan-2-yl)-2-hydroxy-1-((S)-1-methylpropyl)butyl)thiophene-2-sulfonamide;
- 5-chloro-N-((S)-2-hydroxy-1-((S)-1-methylpropyl)hex-5-enyl)thiophene-2-sulfonamide;
- 15 5-chloro-N-((S,S)-1-{hydroxy[4-(methylthio)phenyl]methyl}-2-methylbutyl)thiophene-2-sulfonamide;
- 5-chloro-N-((S,S)-1-[[4-(dimethylamino)phenyl](hydroxy)methyl]-2-methylbutyl)thiophene-2-sulfonamide;
- N-((S,S)-1-[cyclopentyl(hydroxy)methyl]-2-methylbutyl)thiophene-2-sulfonamide;
- 20 N-((S)-2-hydroxy-1-((S)-1-methylpropyl)octyl)thiophene-2-sulfonamide;
- N-((S)-2-hydroxy-1-((S)-1-methylpropyl)heptyl)thiophene-2-sulfonamide;
- 25 N-((S)-2-hydroxy-1-((S)-1-methylpropyl)hexyl)thiophene-2-sulfonamide;
- N-((S,S)-1-[hydroxy(2-methylphenyl)methyl]-2-methylbutyl)thiophene-2-sulfonamide;
- N-((S)-2-hydroxy-3,3-dimethyl-1-((S)-1-methylpropyl)butyl)thiophene-2-sulfonamide;
- 30 N-((S)-2-hydroxy-1-((S)-1-methylpropyl)but-3-enyl)thiophene-2-sulfonamide;

N-{(S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl} thiophene-2-sulfonamide;

N-{(S)-2-hydroxy-1-[(S)-1-methylpropyl]butyl} thiophene-2-sulfonamide;

5 N-{(S,S)-1-[hydroxy(4-methoxyphenyl)methyl]-2-methylbutyl} thiophene-2-sulfonamide;

N-{(S)-4-(1,3-dioxan-2-yl)-2-hydroxy-1-[(S)-1-methylpropyl]butyl} thiophene-2-sulfonamide;

10 N-{(S)-2-hydroxy-1-[(S)-1-methylpropyl]hex-5-enyl} thiophene-2-sulfonamide;

N-{(S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-3-ynyl} thiophene-2-sulfonamide;

N-((S,S)-1-{hydroxy[4-(methylthio)phenyl]methyl}-2-methylbutyl)thiophene-2-sulfonamide;

15 N-{(S,S)-1-[[4-(dimethylamino)phenyl](hydroxy)methyl]-2-methylbutyl} thiophene-2-sulfonamide;

5-chloro-N-{(S,S)-1-[(S)-cyclohex-2-en-1-yl(hydroxy)methyl]-2-methylbutyl} thiophene-2-sulfonamide;

20 5-chloro-N-{(S,S,E)-2-hydroxy-1-[(S)-1-methylpropyl]hex-4-enyl} thiophene-2-sulfonamide;

5-chloro-N-{(S,R,E)-2-hydroxy-1-[(S)-1-methylpropyl]hex-4-enyl} thiophene-2-sulfonamide;

5-chloro-N-{(S,R,E)-2-hydroxy-1-[(S)-1-methylpropyl]hept-4-enyl} thiophene-2-sulfonamide;

25 5-chloro-N-{(S,S)-2-hydroxy-4-methyl-1-[(S)-1-methylpropyl]pent-4-enyl} thiophene-2-sulfonamide;

5-chloro-N-{(S,R)-2-hydroxy-4-methyl-1-[(S)-1-methylpropyl]pent-4-enyl} thiophene-2-sulfonamide;

30 5-chloro-N-{(S,E)-2-hydroxy-1-[(S)-1-methylpropyl]-5-phenylpent-4-enyl} thiophene-2-sulfonamide;

5-chloro-N-[(S,S)-1-(1-hydroxy-1-methylethyl)-2-methylbutyl]thiophene-2-sulfonamide;

- 5-chloro-N-((S)-2-hydroxy-1-[(S)-1-methylpropyl]-2-pentylheptyl)thiophene-2-sulfonamide;
- 5-chloro-N-((S,S)-1-[hydroxy(diphenyl)methyl]-2-methylbutyl)thiophene-2-sulfonamide;
- 5 N-((S)-2-allyl-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl)-5-chlorothiophene-2-sulfonamide;
- 5-chloro-N-((S)-2-ethyl-2-hydroxy-1-[(S)-1-methylpropyl]butyl)thiophene-2-sulfonamide;
- 10 N-((S,S)-1-[bis(4-chlorophenyl)(hydroxy)methyl]-2-methylbutyl)-5-chlorothiophene-2-sulfonamide;
- 5-chloro-N-((S)-2-hydroxy-2-isopropenyl-3-methyl-1-[(S)-1-methylpropyl]but-3-enyl)thiophene-2-sulfonamide;
- 5-chloro-N-((S,S)-1-{hydroxy[bis(4-methoxyphenyl)]methyl}-2-methylbutyl)thiophene-2-sulfonamide;
- 15 5-chloro-N-((S,E)-2-hydroxy-3-methyl-2-[(E)-1-methylprop-1-enyl]-1-[(S)-1-methylpropyl]pent-3-enyl)thiophene-2-sulfonamide;
- N-((S)-2-but-3-enyl-2-hydroxy-1-[(S)-1-methylpropyl]hex-5-enyl)-5-chlorothiophene-2-sulfonamide;
- 5-chloro-N-((S,S)-1-{hydroxy[di(1-naphthyl)]methyl}-2-methylbutyl)thiophene-2-sulfonamide;
- 20 5-bromo-N-((S)-2-ethyl-2-hydroxy-1-[(S)-1-methylpropyl]butyl)thiophene-2-sulfonamide;
- 5-bromo-N-((S)-2-hydroxy-2-isopropenyl-3-methyl-1-[(S)-1-methylpropyl]but-3-enyl)thiophene-2-sulfonamide;
- 25 5-bromo-N-((S,E)-2-hydroxy-3-methyl-2-[(E)-1-methylprop-1-enyl]-1-[(S)-1-methylpropyl]pent-3-enyl)thiophene-2-sulfonamide;
- 5-bromo-N-((S)-2-but-3-enyl-2-hydroxy-1-[(S)-1-methylpropyl]hex-5-enyl)thiophene-2-sulfonamide;
- 5-chloro-N-[1-(hydroxymethyl)cyclohexyl]thiophene-2-sulfonamide;
- 30 5-chloro-N-[2-(hydroxymethyl)bicyclo[2.2.1]hept-2-yl]thiophene-2-sulfonamide;

5-chloro-N-[1-(hydroxymethyl)-2,3-dihydro-H-inden-1-yl]thiophene-2-sulfonamide;

5-chloro-N-[2-(hydroxymethyl)-2,3-dihydro-H-inden-2-yl]thiophene-2-sulfonamide;

5 5-bromo-N-[1-(hydroxymethyl)cyclohexyl]thiophene-2-sulfonamide;

5-bromo-N-[2-(hydroxymethyl)bicyclo[2.2.1]hept-2-yl]thiophene-2-sulfonamide;

5-bromo-N-[2-(hydroxymethyl)-2,3-dihydro-H-inden-2-yl]thiophene-2-sulfonamide;

10 5-chloro-N-[(S,S)-1-[(S)-1-hydroxyethyl]-2-methylbutyl]thiophene-2-sulfonamide;

5-chloro-N-[(S,S)-1-[R-1-hydroxyethyl]-2-methylbutyl]thiophene-2-sulfonamide;

15 5-chloro-N-[(S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl]thiophene-2-sulfonamide;

5-chloro-N-[(S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl]thiophene-2-sulfonamide;

5-chloro-N-[(S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl]thiophene-2-sulfonamide;

20 5-chloro-N-[(S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl]thiophene-2-sulfonamide;

5-bromo-N-[(S,S)-1-[(S)-1-hydroxyethyl]-2-methylbutyl]thiophene-2-sulfonamide;

25 5-bromo-N-[(S,S)-1-[R-1-hydroxyethyl]-2-methylbutyl]thiophene-2-sulfonamide;

5-bromo-N-[(S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl]thiophene-2-sulfonamide;

5-bromo-N-[(S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pentyl]thiophene-2-sulfonamide;

30 5-bromo-N-[(S,S)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl]thiophene-2-sulfonamide;

- 5-bromo-N-[(S,R)-2-hydroxy-1-[(S)-1-methylpropyl]pent-4-enyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(S,S)-2-methyl-1-(2,2,2-trifluoro-1-hydroxyethyl)butyl]thiophene-2-sulfonamide;
- 5 5-chloro-N-[1-(1-hydroxybut-3-enyl)cyclohexyl]thiophene-2-sulfonamide;
- 5-chloro-N-[1-(1-hydroxy-3-methylbut-3-enyl)cyclohexyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(S)-2-hydroxy-1-(4-methoxycyclohexyl)ethyl]thiophene-2-sulfonamide;
- 10 5-chloro-N-[(S)-2-hydroxy-1-(4-propoxycyclohexyl)ethyl]thiophene-2-sulfonamide;
- N-[(S)-1-[4-(allyloxy)cyclohexyl]-2-hydroxyethyl]-5-chlorothiophene-2-sulfonamide;
- 15 N-[(S)-1-[4-(benzyloxy)cyclohexyl]-2-hydroxyethyl]-5-chlorothiophene-2-sulfonamide;
- N-[1-acetyl-4-(hydroxymethyl)piperidin-4-yl]-5-chlorothiophene-2-sulfonamide;
- N-[(1S)-2-butyl-1-(hydroxymethyl)hexyl]-5-chloro-2-thiophenesulfonamide;
- 20 N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-5-iodo-2-thiophenesulfonamide;
- 5-fluoro-N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]-2-thiophenesulfonamide;
- 25 N-[(1S,2S)-1-(hydroxymethyl)-2-methylbutyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(S)-2-hydroxy-1-(4-benzylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 5-chloro-N-[(S)-2-hydroxy-1-(4-methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 30 5-chloro-N-[(S)-2-hydroxy-1-(4-ethylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;

- 5-chloro-*N*-[(*S*)-2-hydroxy-1-(4-npropylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 5-chloro-*N*-[(*S*)-2-hydroxy-1-(4-allylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 5 5-chloro-*N*-[(*S*)-2-hydroxy-1-(4-(3-pyridyl)methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 5-chloro-*N*-[(*S*)-2-hydroxy-1-(4-morpholinocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 5-chloro-*N*-[(*S*)-2-hydroxy-1-(4-(4-pyridyl)methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 10 5-chloro-*N*-[(*S*)-2-hydroxy-1-(4-(2-pyridyl)methylaminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 5-chloro-*N*-[(*S*)-2-hydroxy-1-(4-(carboethoxymethyl)aminocyclohexyl)ethyl]thiophene-2-sulfonamide;
- 15 5-Chloro-*N*-[(*S*)-2-hydroxy-1-(4-hydroxycyclohexyl)ethyl]thiophene-2-sulfonamide
- 5-chloro-*N*-[(*S*)-2-ethyl-1-formylbutyl]thiophene-2-sulfonamide;
- 5-chloro-*N*-[(*S*)-2-ethyl-1-(1-hydroxyethyl)butyl]thiophene-2-sulfonamide;
- 20 5-chloro-*N*-[(*S*)-2-ethyl-1-(1-hydroxy-1-methylethyl)butyl]thiophene-2-sulfonamide;
- 5-chloro-*N*-(2-hydroxy-1-tetrahydro-*H*-thiopyran-4-ylethyl)thiophene-2-sulfonamide;
- 5-chloro-*N*-[(*S*)-2-hydroxy-1-piperidin-4-ylethyl]thiophene-2-sulfonamide;
- 25 *N*-[(*S*)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide;
- N*-[(*S*)-2-ethyl-1-(hydroxymethyl)butyl]-5-fluorothiophene-2-sulfonamide;
- 5-chloro-*N*-[(*S*)-1-(2,3-dihydro-*H*-inden-2-yl)-2-hydroxyethyl]thiophene-2-sulfonamide;
- 30 5-chloro-*N*-{(*S,S*)-1-[(*Z*)-(hydroxyimino)methyl]-2-methylbutyl}thiophene-2-sulfonamide; and

5-chloro-*N*-{(*S,S*)-1-[(*E*)-(hydroxyimino)methyl]-2-methylbutyl}thiophene-2-sulfonamide.

18. The compound according to claim 1, wherein the compound is 5-chloro-*N*-[(1*S*)-2-ethyl-1-(hydroxymethyl)butyl]thiophene-2-sulfonamide.
19. The compound according to claim 1 or 5, wherein X is O, and W, Y and Z are independently selected from C and CR₁₀.
20. The compound according to claim 19, wherein R₆ is halogen, R₄ is lower alkyl of *S*-stereochemistry, R₃ is hydrogen, R₅ is hydrogen, and R₁ and R₂ are each hydrogen.
21. The compound according to claim 1, wherein the compound is selected from the group consisting of *N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide and 5-chloro-*N*-[(1*S*,2*S*)-1-(hydroxymethyl)-2-methylbutyl]-2-furansulfonamide.
22. The compound according to claim 1, wherein the compound is 4-[1-(5-chloro-thiophene-2-sulfonylamino)-2-hydroxy-ethyl]-piperidine-1-carboxylic acid tert-butyl ester.
23. The compound according to any of claims 1 to 22, wherein the pharmaceutically acceptable salt is selected from the group consisting of salts of acetic acid, lactic acid, citric acid, tartaric acid, succinic acid, fumaric acid, maleic acid, malonic acid, mandelic acid, malic acid, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, methanesulfonic acid, toluenesulfonic acid, salts of bases, and mixtures thereof.
24. A pharmaceutical composition comprising a compound according to any of claims 1 to 23 and a physiologically compatible carrier.

25. A method of inhibiting beta amyloid production in a subject, said method comprising the step of delivering a compound according to any of claims 1 to 23.

5 26. The method according to claim 25, wherein said compound is delivered orally or by injection.

10 27. A method of treating a disease selected from the group consisting of Alzheimer's Disease, amyloid angiopathy, cerebral amyloid angiopathy, systemic amyloidosis, hereditary cerebral hemorrhage with amyloidosis of the Dutch type, inclusion body myositis, and Down's syndrome, in a subject, said method comprising the step of administering a compound according to any of claims 1 to 23 to said subject in an amount sufficient to alleviate the symptoms or progress of said disease.

15 28. Use of the compound according to any of claims 1 to 23, or said acceptable salt thereof, in preparing a medicament.

20 29. Use of the compound according to any of claims 1 to 23, or said acceptable salt, in preparing a medicament useful for administration to a mammalian subject for inhibiting beta amyloid production.

30. The use according to claim 29, wherein said compound is administered orally or by injection.

25 31. Use of the compound according to any of claims 1 to 23, or said acceptable salt, in preparing a medicament useful for administration to a mammalian subject for treating a disease selected from the group consisting of Alzheimer's Disease, amyloid angiopathy, cerebral amyloid angiopathy, systemic amyloidosis, hereditary cerebral hemorrhage with amyloidosis of the Dutch type, inclusion body myositis, and Down's syndrome, wherein said compound is administered in an amount sufficient to alleviate the symptoms or progress of said disease.

30